



Twin-singleton differences in intelligence?

Daniëlle Posthuma¹, Eco JC De Geus¹, Nico Bleichrodt² and Dorret I Boomsma¹

¹Department of Biological Psychology and ²Department of Work and Organizational Psychology, Vrije Universiteit Amsterdam, The Netherlands

The twin method has been criticised for its alleged non-generalisability. When population parameters of intellectual abilities are estimated from a twin sample, critics point to the twin-singleton differences in intrauterine and family environments. These differences are suggested to lead to suboptimal cognitive development in twins. Although previous studies have reported twin-singleton differences in intelligence, these studies had two major drawbacks: they tested young twins, and twins were compared with (genetically) unrelated singletons. To test accurately whether twin-singleton differences in intelligence exist, a group of adult twins and their non-twin siblings were administered the Dutch WAIS-III. The group was large enough to detect twin-singleton differences of magnitudes reported in earlier investigations. The data were analysed using maximum likelihood model fitting. No evidence of differences between adult twins and their non-twin siblings on cognitive performance was found. It is concluded that twin studies provide reliable estimates of heritabilities of intellectual abilities which can be generalised to the singleton population. *Twin Research* (2000) 3, 83–87.

Keywords: twin study, intelligence, twins, singletons

Introduction

Classic behavioural genetic studies provide statistical estimates of heritabilities that form the first step in the search for genes for complex behaviour.^{1,2} A large part of these behavioural genetic studies are based on twin samples. These samples have sometimes been criticised for their alleged non-generalisability; since twins are 'special' they may not be representative of singletons. Especially in the field of cognitive abilities twins are generally considered to be at a disadvantage compared with singletons.^{3–6}

Twins share the womb at the same time and consequently share prenatal nutrition provided by the mother's dietary intake. When preparing for labour, twins compete for the best position. This suboptimal intrauterine environment may lead to prematurity, low birth weight and lower weight-for-gestational age,⁷ which in turn in several cases have been associated with low childhood IQ.^{8–12} Apart from a general suboptimal intrauterine environment for both twins, it is known that one of the two foetuses will suffer more from this suboptimal environment than the other.¹³ It is usually the second-born twin that experiences the greatest adverse effects of sharing the womb.¹⁴

Beside these adverse effects of sharing the womb twins may suffer from twin-related stresses in the

family environment in which they are reared. A multiple birth puts stress on a family which may have a negative effect on the (cognitive) development of a twin pair. In some studies it is argued that especially for monozygotic (MZ) twins, who are very much alike, limitation of resources and competition may lead to negative influences for at least one twin member.³

A relatively small number of studies has been devoted to detecting twin-singleton differences in cognition.^{4,6,15} The one study that stands out was conducted by Record, McKeown and Edwards⁶ who compared an impressive number of singletons, twins and even a few triplets. Verbal reasoning scores from the British eleven-plus examination were gathered from 48 913 singletons, 1082 twin pairs and eleven triplets. Standard verbal reasoning scores were significantly lower for twins (standard verbal IQ 95.7) than for singletons (100.1). Triplets performed even worse (91.6). The authors investigated whether this 4.4 standard points difference between twins and singletons could be attributed to effects of maternal age, birth weight, gestational age, zygosity and whether a twin was born first or second. None of these factors could explain the difference.

Record *et al*⁶ also investigated whether twins of whom one had died shortly after birth differed from singletons; although for these 'twins' a slightly lower score than normal singletons (1.9 points) was found, this difference was much smaller than the 4.4 points difference between singletons and twins of which both members were still alive. Based on this observation the authors concluded that the difference of

Correspondence: D Posthuma, Vrije Universiteit, Department of Biological Psychology, De Boelelaan 1111, 1081 HV, Amsterdam, The Netherlands. Tel: +31 20 444 8814; Fax: +31 20 444 8832; E-mail: danielle@psy.vu.nl

Received 15 June 1999; accepted 15 June 1999

4.4 points between singletons and twins cannot be attributed to negative effects of sharing the womb, but instead must be sought in the environment in which twins are reared. However, since Record *et al*⁶ did not control for any difference in twin families and singleton families, they could not rule out selection biases in the sampling of twin and non-twin families. Such biases may exist because twins as a group may have a slightly different genetic or social background than singletons.

Nathan and Guttman¹⁶ tried to overcome selection bias in twin and singleton families by comparing twins and singletons (aged 8–13 years) who were reared in the same kibbutz. A kibbutz is an Israeli community in which children are collectively reared. So although the twins and singletons in this study did not have the same genetic background, they were accurately matched for family environment and childrearing practices. In this study dizygotic (DZ) twins performed worse than MZ twins and singletons. According to the authors, however, this difference could be totally ascribed to the relatively few years of schooling of the group of DZ mothers. Thus, in spite of the attempt to match twins and singletons this study is also an example of biased family sampling.

In addition to comparing twins with familiarly unrelated singletons, most previous studies have been conducted using young twins.^{9,10,17–20} Because these studies show that twins recover any deficits in intellectual performance by 6–8 years of age,^{18–20} the comparison of twins and singletons at ages below 8 years does not provide a good indication of adult twin–singleton differences. To the best of our knowledge studies comparing the IQ of adult twins and genetically related singletons have not yet been conducted.

In the present study mean scores of adult MZ and DZ twins on intellectual ability are compared with the mean scores of their non-twin siblings. Non-twin siblings make an ideal control group; both genetic background and early familial environments are perfectly matched.

Method

Subjects

The subjects were 358 family members from a total of 152 twin families who participated in a project investigating the genetics of adult brain function. The Dutch version of the Wechsler Adult Intelligence Scale-III (WAIS-III)²¹ was administered when the participants visited the laboratory for a combined session of neuropsychological and electroencephalographic measurements. All subjects were recruited from the Netherlands Twin Registry. The

twins had previously participated in one of two previously conducted studies in which zygosity was assessed by blood group polymorphisms and DNA typing.^{22,23}

In total, 98 siblings, 101 MZ twins, 153 DZ twins and 9 triplets participated. Since the group of triplets was small, we discarded the data of the last born of the triplets and treated the remaining two members as if they were twins. This left 98 siblings and 260 twins. The study recruited twin pairs and at most two of their non-twin siblings. It also included single twins (co-twin refused participation) and siblings only (both twins refused). Thus, families consisted of at least one member and at most four members. Table 1 shows the number of families with a particular constitution, eg 27 MZ families consisting of two twin members and no siblings participated; siblings from nine families participated without the twins. Due to administrative errors five individual test scores are missing subtest *digit symbol-coding*, four individual test scores are missing subtests *block design* and *digit symbol-free recall*, and one individual test score is missing subtest *digit symbol-pairing* and subtest *letter-number sequencing*. Results are based on the available number of subjects per subtest (see Table 3).

Mean age and sex distribution per group are displayed in Table 2. Of the 98 non-twin siblings, 35 were younger than the twin from the same family, and 63 were older. Distribution of sex did not differ in the DZ twins and the siblings. Slightly fewer female MZ twins than male MZ twins participated.

Table 1 Sample configuration

		number of non-twin siblings			
		0	1	2	
mz twins	2 twins	27	18	3	total mz twin pairs: 48
	1 twin	–	4	1	
dz twins	2 twins	32	27	10	total dz twin pairs: 69
	1 twin	12	8	1	
no twin		–	7	2	
total non-twin siblings:		64	+ 34	= 98	

Table 2 Mean age and sex distribution per group

Group	Male	Female	Total	Mean age in years (sd)
mz twins	58	43	101	39.7 (12.63)
dz twins	70	89	159	37.3 (11.87)
sibs	46	52	98	37.1 (12.02)

sd = standard deviation

Procedure

Eleven subtests of the Dutch WAIS-III were administered in a fixed order. Subtests included *block design, letter-number sequencing, information, matrix reasoning, similarities, picture completion, arithmetic, vocabulary, digit symbol coding, digit symbol pairing and digit symbol free recall*. Age and sex normalised scores for the Dutch WAIS-III are not yet available; raw scores were used in the analyses throughout. All subjects were paid Dfl. 50.- for participation.

Statistical analyses

As can be seen from Table 1 the data were characterised by the varying number of participating family members; families consisted of one to four members which could be any combination of one or two twins and/or non-twin siblings. This variability in number of observations per family causes serious computational problems. In Mx²⁴ the handling of such 'incomplete' data is implemented by calculating twice the negative log-likelihood (-LL) of the raw data for each family, with the following formula:

$$-LL = -k \log (2\pi) + \log |\Sigma| + (x_i - \mu_i)' \Sigma^{-1} (x_i - \mu_i),$$

where k ($k = 1, 2, 3$ or 4) denotes the number of observed variables within a family, Σ (4×4) is the covariance matrix of family members, x_i (for $i = 1, 2, 3, 4$) is the vector of observed scores, μ_i is the column vector of the estimated means of the variables, and $|\Sigma|$ and Σ^{-1} are the determinant and inverse of matrix Σ , respectively.

When two models which provide -2LLs are nested, subtracting the two -2LLs from each other provides a $\Delta(-2LL)$ which has a χ^2 distribution. A high χ^2 against a low gain of degrees of freedom (Δdf) denotes a worse fit of the second, more restrictive model relative to the first model.

Four univariate nested models were fitted using this procedure. In the first model all means were estimated individually. The second model is the same as the first model with two extra equality constraints; one on the means of both members of the MZ twin pairs and another one on the means of both members of the DZ twin pairs. The third model is the same as the second model but further constrains the means of the MZ twin pairs and the DZ twin pairs to be equal. The fourth is the same as the third model but with an extra equality constraint on the means of all twins (mz and dz) and siblings.

Model 2 tests whether the means of first born twins and second born twins within zygosity groups are significantly different. The third model serves as a test of the assumption that the means in MZ twins

and DZ twins do not differ. Model 4 tests whether the means of twins and siblings are significantly different.

For all models the variances of all twin members and all siblings were constrained equal, and all covariances of all twin sib pairs, the covariance of two sibs within one family and the covariance of the DZ twins were set equal.

Statistical power

We calculated the necessary sample size for each group (singletons and twins) based on the effect size as found in Record *et al's* study.⁶ A measure of effect size that is independent of scaling is Cohen's d , which is calculated as follows:

$$d = (\mu_1 - \mu_2)/\sigma$$

where μ_1 is the mean of the first group (singletons), μ_2 is the mean of the second group (twins) and σ is the common standard deviation.²⁵

Record *et al*⁶ found a 4.4 standard points difference between the two groups. The standard deviation of an IQ score is by definition 15. The effect size in the Record *et al* study was thus 0.29, which is considered a small effect. For a one-tailed test with $\alpha = 0.05$, $1 - \beta = 0.80$, and two related samples, 70 individuals per group (singletons and twins) are needed to detect an effect of such small magnitude.²⁶ We had 260 twins and 98 non-twin siblings giving us the power to detect effect sizes well below 0.29.

Results

The observed means and standard deviations of WAIS-III subtests per group are displayed in Table 3.

Table 3 Observed means and standard deviations of WAIS-III subtests per group

subtest	mz twin (N = 101)	dz twin (N = 159)	sibs (N = 98)
Block design	26.20 (8.96)	25.72 (9.28) ^a	26.25 (8.85) ^b
LN sequencing	12.21 (3.42)	11.21 (2.61)	11.86 (2.90) ^c
Information	23.41 (6.32)	23.93 (6.00)	24.11 (6.54)
Matrix reasoning	19.36 (3.38)	19.16 (3.44)	19.40 (3.28)
Similarities	26.91 (5.58)	27.17 (5.43)	27.33 (5.58)
Picture completion	20.86 (2.55)	20.72 (2.60)	20.55 (3.18)
Arithmetic	13.86 (3.86)	13.75 (3.89)	14.70 (4.12)
Vocabulary	49.07 (11.60)	48.26 (10.55)	47.83 (13.54)
DS coding	76.09 (15.22)	77.66 (19.52) ^d	78.83 (15.86) ^e
DS free recall	7.63 (1.20) ^f	7.54 (1.12) ^d	7.54 (1.27) ^c
DS pairing	13.25 (4.25)	12.67 (4.19)	12.92 (4.02) ^c

^abased on 157 observations

^bbased on 96 observations

^cbased on 97 observations

LN = Letter-number

^dbased on 158 observations

^ebased on 94 observations

^fbased on 99 observations

DS = Digit symbol

Table 4 Fit indices for nested sequence of models fitted to raw data of WAIS-III subtest scores of MZ twins, DZ twins and siblings

Subtest	1. All means unequal		2. Means 1st born twins equal means 2nd born twins, within zygosity groups		3. Means mz twins equal means dz twins		4. Means twins equal means non-twin siblings		(4-1) All means equal against all means unequal	
	-2LL	df	-2LL	df	-2LL	df	-2LL	df	χ^2	($\Delta df = 7$) ^a
Block design	2451.48	343	2453.90	345	2454.01	346	2459.56	350	8.08	n.s.
Letter-number sequencing	1738.22	346	1739.38	348	1744.57	349	1750.37	353	12.15	n.s.
Information	2194.37	347	2197.44	349	2197.87	349	2205.64	354	11.27	n.s.
Matrix reasoning	1842.22	347	1845.75	349	1845.93	350	1848.00	354	5.78	n.s.
Similarities	2150.00	347	2151.07	349	2151.21	350	2157.71	354	7.71	n.s.
Incomplete pictures	1681.34	347	1681.81	349	1681.85	350	1687.18	354	5.84	n.s.
Arithmetic	1919.46	347	1920.33	349	1920.44	350	1930.52	354	11.06	n.s.
Vocabulary	2675.27	347	2678.30	349	2678.60	350	2682.41	354	7.14	n.s.
Digit symbol coding	2964.08	342	2965.69	344	2965.99	345	2967.20	349	3.12	n.s.
Digit symbol free recall	1082.13	343	1082.29	345	1082.61	346	1092.84	350	10.71	n.s.
Digit symbol pairing	1988.00	346	1990.40	348	1991.25	349	1994.45	353	6.45	n.s.

df = degrees of freedom; -2LL = twice the negative log likelihood; n.s. = not significant: when the increase in χ^2 is not significant, the most restrictive model is accepted; ^aan increase in χ^2 of more than 14.07 for $\Delta df = 7$ is significant at the 0.05 level.

To test whether the above differences in mean scores indicated true differences, univariate analyses in Mx using twice the negative log-likelihood were run. The results for these analyses are presented in Table 4, from which it can be seen that comparison of model 4, the most parsimonious model, with model 1 did not cause a significant worsening of the fit for any of the WAIS III subtests. In other words, for all subtests a model which estimates all means to be equal fits better than a model in which all means are estimated separately. There was no reason to believe that means of twins and singletons in our sample differed in IQ.

We did find, however, that comparison of model 4 (all means equal) with model 3 (separate means for twins and siblings) showed a significant worsening of the fit for subtests *arithmetic* and *digit symbol-free recall*, in the sense that on *arithmetic* singletons performed slightly better than both MZ and DZ twins, and on *digit symbol-free recall* MZ twins performed slightly better than both DZ twins and singletons. We also found that MZ twins performed significantly better than DZ twins on subtest *letter-number sequencing*.

Discussion

It has been suggested that twins have an intellectual disadvantage compared with singletons and that twin samples are not representative of the normal population. If true, this might influence generalisability of heritability estimates obtained in twin studies, for instance by a restriction of range of IQ scores. In the Record *et al*⁶ study a standard IQ score difference of 4.4 points was found between twins and singletons. Our study had enough statistical power to detect an effect of at least the same magnitude on each of the individual IQ subtests. We

found, however, no evidence of a twin-singleton difference. In fact, means and standard deviations in our study showed no differences at all between twins and singletons. In the Record *et al*⁶ study, where these differences were found, *a priori* differences in social class or genetic background of twin families and singleton families could never be ruled out. Since our twins and singletons came from the same family, social class and genetic background were perfectly matched across twin families and singleton families.

Our results are in line with an earlier report by Kallman²⁷ who administered the Wechsler Bellevue Scale to 134 twin pairs (aged 60–89 years), and compared the scores of these twins to standardised scores based on a comparable group of singletons. Kallman concluded that there was no significant difference between twins and singletons in measures of intellectual performance.

Although in our study no evidence was found for twin-singleton differences in intellectual ability, one cannot necessarily generalise from this in respect of personality, lifestyle, disease susceptibility or mortality rates. However, recent comparisons of twins and singletons on problem behaviour,²⁸ mortality rates²⁹ and psychiatric symptoms³⁰ have not suggested twin-singleton differences in these fields either. All in all, significant disadvantages of twins in comparison with singletons seem to be implied rather than observed.

Acknowledgements

We gratefully acknowledge the financial support of the USF (grant number 96/22) and the HFSP (grant number rg0154/1998-B).

References

- 1 Bouchard TJ, Lykken DT, McGue M, Segal NL, Tellegen A. Sources of human psychological differences: The Minnesota study of twins reared apart. *Science* 1990; **250**: 223–228.
- 2 McClearn GE, Johansson B, Berg S, Pedersen NL, Ahern F, Petrill S, Plomin R. Substantial genetic influence on cognitive abilities in twins 80 or more years old. *Science* 1997; **276**: 1560–1563.
- 3 Hay DA, O'Brien PJ. The La Trobe Twin Study: A genetic approach to the structure and development of cognition in twin children. *Child Devel* 1983; **54**: 317–330.
- 4 Hay DA, O'Brien PJ, Johnston CJ, Prior MR. The high incidence of reading disability in twin boys and its implications for genetic analyses. *Acta Genet Med Gemellol Roma* 1984; **33**: 223–236.
- 5 Vandenberg SG. Does a special twin situation contribute to similarity for abilities in MZ and DZ twins? *Acta Genet Med Gemellol Roma* 1984; **33**: 219–222.
- 6 Record RG, McKeown T, Edwards JH. An investigation of the difference in measured intelligence between twins and single births. *Ann Hum Genet* 1970; **34**: 11–20.
- 7 Powers WF, Kiely JL. The risks confronting twins: A national perspective. *Am J Obstet Gynaecol* 1994; **170**: 456–461.
- 8 Record RG, McKeown T, Edwards JH. The relation of measured intelligence to birth weight and duration of gestation. *Ann Hum Genet* 1969; **33**: 71–79.
- 9 Alin-Akerman B. Eight-year follow-up of cognitive development in 33 twin pairs. *Acta Genet Med Gemellol Roma* 1995; **44**: 179–188.
- 10 Myriantopoulos NC, Nichols PL, Broman SH. Intellectual development of twins comparison with singletons. *Acta Genet Med Gemellol Roma* 1976; **25**: 376–380.
- 11 Willerman L, Churchill JA. Intelligence and birth weight in identical twins. *Child Devel* 1967; **38**: 623–629.
- 12 Lynn R, Hattori K. The heritability of intelligence in Japan. *Behav Genet* 1990; **20**: 545–546.
- 13 Price B. Primary biases in twin studies. *Am J Hum Genet* 1950; **2**: 293–352.
- 14 Prins RP. The second born twin: can we improve outcomes? *Am J Obstet Gynaecol* 1994; **170**: 1649–1657.
- 15 Lange AL, Fischbein S. From puberty to mid-life: a follow-up study of twins and controls. *Acta Genet Med Gemellol Roma* 1992; **41**: 105–112.
- 16 Nathan M, Guttman R. Similarities in test scores and profiles of kibbutz twins and singletons. *Acta Genet Med Gemellol Roma* 1984; **33**: 213–218.
- 17 Dezoete JA, MacArthur BA. Cognitive development and behaviour in very low birthweight twins at four years. *Acta Genet Med Gemellol Roma* 1996; **45**: 325–332.
- 18 Wilson RS. Twin growth: initial deficit, recovery, and trends in concordance from birth to nine years. *Ann Hum Biol* 1979; **6**: 205–220.
- 19 Wilson RS. Twins: Patterns of cognitive development as measured on the Wechsler Pre-school and Primary Scale of Intelligence. *Development Psychol* 1975; **11**: 126–134.
- 20 Wilson RS. Twins: Mental development in the preschool years. *Development Psychol* 1974; **10**: 580–588.
- 21 *WAIS-III Manual* (Dutch version) Swets and Zeitlinger: Lisse, 1997.
- 22 Boomsma DI, Kempen HJM, Gevers-Leuven JA, Havekes L, de Knijff P, Frants RR. Genetic analysis of sex and generation differences in plasma lipid, lipoprotein and apolipoprotein levels in adolescent twins and their parents. *Genet Epidemiol* 1996; **13**: 49–60.
- 23 Snieder H, Doornen LJP van, Boomsma DI. Age-dependency of gene expression for plasma lipids, lipoproteins and apolipoproteins. *Am J Hum Genet* 1997; **60**: 638–650.
- 24 Neale MC. *Mx: Statistical modeling*. 3rd edn. MCV: Richmond VA, 1997.
- 25 Cohen J. A power primer. *Psychol Bull* 1992; **112**: 155–159.
- 26 Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd edn. Lawrence Erlbaum Associates: New Jersey, 1988.
- 27 Kallman FJ, Feingold L, Bondy E. Comparative adaptational, social, and psychometric data on the life histories of senescent twin pairs. *Am J Hum Genet* 1951; **3**: 65–73.
- 28 Oord EJCG van den, Koot HM, Boomsma DI, Verhulst FC, Orlebeke JF. A twin-singleton comparison of problem behavior in 2–3 year olds. *J Child Psychol Psychiatr* 1995; **36**: 449–458.
- 29 Christensen K, Vaupel JW, Holm NV, Yashin AI. Mortality among twins after age 6: fetal origins hypothesis versus twin method. *BMJ* 1995; **310**: 432–436.
- 30 Kendler KS, Martin NG, Heath AC, Eaves LJ. Self-report psychiatric symptoms in twins and their nontwin relatives: Are twins different? *Am J Med Genet* 1995; **60**: 588–591.