

## ORIGINAL ARTICLE

# Poor replication of candidate genes for major depressive disorder using genome-wide association data

FJ Bosker<sup>1,8</sup>, CA Hartman<sup>1,8</sup>, IM Nolte<sup>2,8</sup>, BP Prins<sup>2</sup>, P Terpstra<sup>2</sup>, D Posthuma<sup>3,4</sup>, T van Veen<sup>5</sup>, G Willemsen<sup>3</sup>, RH DeRijk<sup>5</sup>, EJ de Geus<sup>3</sup>, WJ Hoogendijk<sup>6</sup>, PF Sullivan<sup>7</sup>, BW Penninx<sup>1,5,6</sup>, DI Boomsma<sup>3</sup>, H Snieder<sup>2,8</sup> and WA Nolen<sup>1,8</sup>

<sup>1</sup>Department of Psychiatry, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands; <sup>2</sup>Unit of Genetic Epidemiology and Bioinformatics, Department of Epidemiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands; <sup>3</sup>Department of Biological Psychology, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands; <sup>4</sup>Department of Medical and Functional Genomics, VU University Medical Center, Amsterdam, The Netherlands; <sup>5</sup>Department of Psychiatry, Leiden University Medical Center, Leiden, The Netherlands; <sup>6</sup>Department of Psychiatry, VU University Medical Center, Amsterdam, The Netherlands and <sup>7</sup>Department of Genetics, University of North Carolina, Chapel Hill, NC, USA

Data from the Genetic Association Information Network (GAIN) genome-wide association study (GWAS) in major depressive disorder (MDD) were used to explore previously reported candidate gene and single-nucleotide polymorphism (SNP) associations in MDD. A systematic literature search of candidate genes associated with MDD in case-control studies was performed before the results of the GAIN MDD study became available. Measured and imputed candidate SNPs and genes were tested in the GAIN MDD study encompassing 1738 cases and 1802 controls. Imputation was used to increase the number of SNPs from the GWAS and to improve coverage of SNPs in the candidate genes selected. Tests were carried out for individual SNPs and the entire gene using different statistical approaches, with permutation analysis as the final arbiter. In all, 78 papers reporting on 57 genes were identified, from which 92 SNPs could be mapped. In the GAIN MDD study, two SNPs were associated with MDD: *C5orf20* (rs12520799;  $P=0.038$ ; odds ratio (OR) AT=1.10, 95% CI 0.95–1.29; OR TT=1.21, 95% confidence interval (CI) 1.01–1.47) and *NPY* (rs16139;  $P=0.034$ ; OR C allele=0.73, 95% CI 0.55–0.97), constituting a direct replication of previously identified SNPs. At the gene level, *TNF* (rs76917; OR T=1.35, 95% CI 1.13–1.63;  $P=0.0034$ ) was identified as the only gene for which the association with MDD remained significant after correction for multiple testing. For *SLC6A2* (norepinephrine transporter (NET)) significantly more SNPs (19 out of 100;  $P=0.039$ ) than expected were associated while accounting for the linkage disequilibrium (LD) structure. Thus, we found support for involvement in MDD for only four genes. However, given the number of candidate SNPs and genes that were tested, even these significant may well be false positives. The poor replication may point to publication bias and false-positive findings in previous candidate gene studies, and may also be related to heterogeneity of the MDD phenotype as well as contextual genetic or environmental factors.

*Molecular Psychiatry* advance online publication, 30 March 2010; doi:10.1038/mp.2010.38

**Keywords:** candidate genes; genome-wide association study; major depressive disorder; replication; single-nucleotide polymorphisms

## Introduction

Major depressive disorder (MDD) is a multi-factorial disease, with both genetic and environmental factors likely to have a role in its etiology. The Netherlands Study of Depression and Anxiety (NESDA; www.

nesda.nl) and the Netherlands Twin Registry (NTR; www.tweelingenregister.org) took part in the Genetic Association Information Network (GAIN; http://www.fnih.org/GAIN) project to enable a genome-wide association study (GWAS) using a 600K Perlegen chip (Perlegen Sciences, Mountain View, CA, USA).<sup>1</sup> Within the GAIN MDD study,<sup>2</sup> 1862 participants with a diagnosis of MDD and 1860 controls at low liability for MDD were selected for genome-wide genotyping.

A GWAS approach allows a hypothesis-free search for potential new susceptibility genes. The downside of a GWAS is that a strict statistical adjustment for the large number of single-nucleotide polymorphisms

Correspondence: Dr FJ Bosker, Department of Psychiatry, University Medical Center Groningen, University of Groningen, PO Box 30.001, 9700 RB Groningen, The Netherlands.  
E-mail: f.j.bosker@psy.umcg.nl

<sup>8</sup>These authors contributed equally to this work.

Received 3 August 2009; revised 18 February 2010; accepted 22 February 2010

(SNPs; in the GAIN MDD study 435 291 SNPs) is required before associations can be considered significant on a genome-wide level,<sup>3</sup> and replication of such findings in independent cohorts is mandatory to exclude false-positive findings.<sup>4</sup>

Another potential use of a GWAS is to use the results for a large-scale replication study of previous candidate gene studies. Application of such previous knowledge within the context of a GWAS allows less stringent significance thresholds than those for the hypothesis-free GWAS analysis.<sup>5,6</sup> So far, the role of candidate genes in MDD has been the subject of many association studies. Unfortunately, there is little consistency between candidate gene studies for multifactorial diseases such as MDD (see, for example, Hirschhorn *et al.*<sup>7</sup> and Munafo<sup>8</sup>).

In the current study we attempted to replicate significant findings from previous candidate gene studies in MDD. To this end we conducted a systematic review of the literature and selected those genes that were reported to be significantly associated with MDD at least once. The GWAS data from the GAIN MDD study were used to screen all the identified candidate genes in two ways: (1) for association with the specific SNPs reported in the literature; and (2) for association with any of the SNPs (genotyped or imputed) from the Perlegen chip within the identified genes.

## Materials and methods

### *Selection of studies reporting on candidate genes*

To prevent any bias from the results of the GAIN MDD study, we identified candidate genes for MDD before the results from the GAIN MDD study became available. Therefore, a so-called 'enhanced search' was performed in Medline through PubMed on 1 September 2007 using the following search terms: (('genes' (TIAB) NOT Medline(SBI)) OR 'genes' (MeSH terms) OR gene (text word)) OR ('genes' (MeSH terms) OR genes (text word)) OR snp (all fields) OR ('single nucleotide polymorphism' (text word) OR 'polymorphism, single nucleotide' (MeSH terms) OR snps (text word)) OR ('genetic polymorphism' (text word) OR 'polymorphism, genetic' (MeSH terms) OR polymorphism (text word)) OR polymorphisms (all fields) AND ('major depressive disorder' (text word) OR 'depressive disorder, major' (MeSH Terms) OR major depression (text word)) AND 'humans' (MeSH terms). This resulted in 641 hits. We additionally scrutinized the reference list of the meta-analysis of genetic studies on MDD by Lopez-Leon *et al.*<sup>9</sup> that appeared online on 16 October 2007, shortly after the end date of our search, resulting in an additional 39 hits of possibly relevant papers. These researchers used somewhat broader search terms than we did, and their search ran until June 2007; therefore, as a final check, we searched the literature using their search terms for the period June 2007 to September 2007 not covered in their paper. This yielded an additional 110 hits. Of all these papers we


retrieved the abstracts, and if considered relevant, the full paper.

In the next step, we selected all papers fulfilling the following five inclusion criteria: (1) The study had to be a candidate gene case-control association study. Linkage and fine-mapping studies were excluded. (2) The primary diagnosis of the patients in the candidate gene study had to be MDD to enable comparison with the GAIN MDD study. Therefore, we excluded studies involving patients: (i) with a depressive episode in the course of bipolar disorder, (ii) with a primary psychotic disorder such as schizophrenia and a secondary depression, (iii) with a seasonal affective disorder not being MDD with a seasonal pattern, (iv) with a primary anxiety disorder (that is, panic disorder, agoraphobia or social phobia) or obsessive compulsive disorder and a secondary depression and (v) with MDD plus an additional specific comorbid disorder or condition, for example, MDD plus alcoholism. However, we allowed subgroups within MDD, for example, MDD in women or in men, recurrent MDD or early-onset MDD. (3) The sample of a specific study consisted of at least 30 patients with MDD and 30 healthy controls. (4) The findings on the association with MDD of any variant within the candidate gene (either a SNP, a microsatellite marker or a haplotype) had reached a statistical significance at the level of  $P < 0.05$ . (5) Finally, the genetic association had to be with the diagnosis of MDD and not with other aspects such as associated personality features (for example, neuroticism) or factors related to treatment response. This resulted in 78 papers.

In order not to miss potential true-positive findings, we did not exclude candidate genes with associated markers that had low allele frequencies or that deviated from Hardy-Weinberg equilibrium. Two investigators (FB and CH) independently made a selection from the initial list of papers using the above-mentioned criteria. When both reached consensus, the paper was included or excluded; in case of disagreement, consensus was obtained with assistance of two other authors (WN and HS). As a final step, one author (IN) double checked whether all selected papers fulfilled the aforementioned selection criteria 1-5. Figure 1 shows a flowchart of how we retrieved the 78 papers for the present study.

### *Bioinformatic tools*

For many SNPs no reference SNP identification number (rs-id) was given in the original papers, but codes based on position (for example, 677C/T in *MTHFR* or Tyr129Ser in *HTR3B*) or even own codes (for example, s1-s5 in *AVPR1B*) were given. To retrieve rs-ids for these SNPs, we used searches in PubMed or in the SNP database of the National Center for Biotechnology Information (NCBI) (<http://www.ncbi.nlm.nih.gov/SNP/>) using the 'Geneview' option in conjunction with NCBI's MapViewer with the human genome assembly build 37.1 (<http://www.ncbi.nlm.nih.gov/projects/mapview/>). Six SNPs



**Figure 1** Selection procedure of studies reporting on candidate genes. Abbreviations: LL, Lopez-Leon; PM, PubMed.

remained that could not be easily found in this way. We contacted the corresponding authors of the papers and used NCBI's Primer-Blast and SNP-Blast (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) to map these SNPs using the provided primer sequences.

#### Sample

The 1862 MDD cases included in the GAIN MDD study were mainly from NESDA, a longitudinal cohort study designed to be representative of individuals with depressive and/or anxiety disorders<sup>10</sup> and were recruited from mental health-care organizations ( $N=785$ ), primary care ( $N=603$ ) and community

samples ( $N=314$ ). Additional cases came from the NTR ( $N=160$ ). Regardless of recruitment setting, similar inclusion and exclusion criteria were used to select MDD cases. Inclusion criteria were a lifetime diagnosis of MDD according to DSM-IV (Diagnostic and Statistical Manual, Fourth Edition)<sup>11</sup> as diagnosed through the Composite International Diagnostic Interview (CIDI Version 2.1.<sup>12</sup>), age 18–65 years and self-reported western European ancestry. People who were not fluent in Dutch and those with a primary diagnosis of a psychotic disorder, obsessive compulsive disorder, bipolar disorder or severe alcohol or substance use disorder were excluded.

Most of the 1860 control subjects were from the NTR ( $N=1,703$ ) and additional controls from NESDA ( $N=157$ ). Longitudinal phenotyping in NTR included assessment of depressive symptoms (through multiple instruments), anxiety and neuroticism. Inclusion required no report of MDD at any measurement occasion and never scoring high ( $>0.5$  s.d.) on a general factor score based on combined measures of neuroticism, anxiety and depressive symptoms. Finally, controls and their parents were required to have been born in the Netherlands or Western Europe. Only one control per family was selected. NESDA controls had no lifetime diagnosis of MDD or an anxiety disorder as assessed by the CIDI and reported low depressive symptoms at baseline. For more details, see Boomsma *et al.*<sup>2</sup>

#### Genotyping

Perlegen Sciences performed all genotyping according to strict standard operating procedures. DNA samples from cases and controls were randomly assigned to plates, shipped to Perlegen and identified only by barcode. High-density oligonucleotide arrays were used yielding 599 164 SNPs. Eight SNPs with duplicate numbers were deleted and 73 mitochondrial SNPs were removed for later analysis. From the remaining 599 083 SNPs on the Perlegen chip, 435 291 passed quality control tests. A total of 280 subjects were excluded because of various quality control issues, leaving 1738 cases (93.4%) and 1802 controls (96.9%) in the final analysis data set. For more details see Boomsma *et al.*<sup>2</sup> and Sullivan *et al.*<sup>4</sup>

#### Imputation

Not all SNPs selected from the literature were present on the genotyping array. On the basis of the linkage disequilibrium (LD) structure between SNPs we followed an imputation procedure to predict non-genotyped SNPs using the HapMap CEU data (release 22, build 36) as the reference database. The imputation was performed by IMPUTE version 0.3.2 using the default settings and the recommended number 11418 for the effective population size of Caucasians.<sup>13</sup> In this way we extended the genome-wide autosomal SNP data set from 427 049 to 2 467 430 SNPs. For our candidate genes this meant an extension from 851 to 4955 SNPs. However, the quality of the imputation was low for 85 SNPs (SNPTEST *proper\_info*  $<0.5$ ). These SNPs were excluded leaving 4870 SNPs for analysis. No SNP had a minor allele frequency of  $<1\%$ .

#### Association test

The association between MDD and the autosomal SNP data was tested using a frequentist case-control test provided in the software package SNPTEST version 1.1.4 using allele dosages with sex as a covariate to adjust for the slight imbalance in the percentage of females between cases (69.6%) and controls (62%),<sup>2,4</sup> and the 'proper' option to account for the uncertainty of the genotypes that were

imputed.<sup>13</sup> In addition, 7988 genotyped SNPs on the X chromosome were analyzed in PLINK version 1.03<sup>14</sup> using logistic regression with sex as a covariate. SNPs on the Y chromosome ( $n=15$ ) and SNPs mapped to ambiguous locations ( $n=239$ ) were not analyzed.

Odds ratios (ORs) and 95% confidence intervals (CIs) were calculated for the significant candidate SNPs according to the disease model from the original article using counted or estimated numbers for genotyped or imputed SNPs, respectively. To establish true replication, we checked whether the effect was for the same allele and in the same direction.

#### Permutation procedure

To facilitate interpretation of the significance of our findings for each SNP, we calculated three  $P$ -values by permutation: (1) a crude uncorrected significance, (2) a gene-wide significance (corrected for all SNPs in the gene) and (3) an overall significance (corrected for all SNPs in all selected genes). The crude uncorrected  $P$ -value was determined as the fraction of permutations that yielded a  $P$ -value that was smaller than the observed one. This  $P$ -value was determined to validate the permutation procedure as it should be similar to the  $P$ -value calculated by SNPTEST or PLINK. The gene-wide significance of a particular SNP was defined as the fraction of permutations that the  $P$ -value of any SNP in the candidate gene concerned was smaller than the one observed for that SNP. This procedure corrects the  $P$ -value for testing multiple SNPs within a gene. On the basis of the rationale that each selected gene is a candidate for MDD (hypothesis driven), no correction for all SNPs in the entire gene set is necessary. Nonetheless, we also calculated an overall significance (that is, corrected for all SNPs in all candidate genes) by determining an overall  $P$ -value for each SNP as the fraction of permutations for which any of the SNPs in any of the candidate genes had a  $P$ -value smaller than the observed one. For each of the three  $P$ -values we conducted 10 000 permutations. Case-control status was randomly permuted within males and females separately, hence leaving the number of affected males and females intact.

In addition to the three SNP-specific  $P$ -values, we also determined whether the number of SNPs within a candidate gene with an original  $P$ -value of  $<0.05$  based on SNPTEST or PLINK (see above) was higher than expected. For each permutation the number of SNPs in each candidate gene that was significant at  $P<0.05$  was recorded. The fraction of permutations with a higher number of significant SNPs than originally observed determined the significance of the number of significant SNPs of that candidate gene. By permuting case-control status the LD structure of the genes is preserved; that is, the resulting significance is corrected for possible high correlation between SNPs. The number of permutations for this test was 10 000 as well.

## Results

### Literature search

The 78 papers that resulted from our systematic literature search reported 115 statistically significant differences between MDD cases and healthy controls in 57 genes: for 96 SNPs, 7 microsatellite markers (that is, length polymorphisms), 11 haplotypes and one protein polymorphism in the haptoglobin gene<sup>15</sup> (Table 1). Twenty-nine SNPs were identified by an rs-id, whereas 67 SNPs were only specified by a location code, restriction enzyme or author-designed code. For 64 of these 67 SNPs, rs-ids could be obtained, whereas two rs-ids were not found (246G/A in *CCKAR*<sup>16</sup> and -7054C/A in *DRD2*<sup>16</sup>). In addition, for SNP 1463G/A in *TPH2*, others have tried to replicate the association, but the SNP seemed to be nonexistent.<sup>17,18</sup> Thus, we could map 93 SNPs.

For the seven microsatellite markers, no LD data with SNPs are known and hence we cannot determine whether these are covered by any of the available SNPs in the corresponding genes. Among them is the 5-HTTLPR 44-bp deletion in the promoter region of the serotonin transporter gene *SLC6A4* (*SERT*), which has attracted considerable attention in various previous candidate gene studies for MDD.

### Association in the GAIN MDD data with specific SNPs from the literature

Of the 93 selected SNPs in the 57 candidate genes, 61 were either present ( $n=18$ ) on the Perlegen array or could be imputed ( $n=43$ ). Four additional SNPs could be tagged by other available SNPs (Table 2). Two of these were not genotyped in the Centre d'Etude du Polymorphisme Humain, Utah (CEU) population of the HapMap Phase2 (release 22, build 36) project, but were available in the Japanese individuals from Tokyo (JPT) and Chinese Han individuals from Beijing (CHB) populations, in which they showed high LD ( $r^2=0.97$  and  $1.00$ , respectively) with at least one other available SNP. The two other SNPs were genotyped in the CEU population and could be tagged by available SNPs with  $r^2=1$ , but were for unreported reasons not included in the HapMap reference file used for the imputation procedure as provided on the website of the imputation software package IMPUTE ([https://mathgen.stats.ox.ac.uk/impute/impute\\_v0.5.html](https://mathgen.stats.ox.ac.uk/impute/impute_v0.5.html)). A total of 28 SNPs were neither genotyped directly nor imputed nor could be tagged, and hence were not available for replication.

We investigated which of the 65 available or tagged SNPs could be confirmed in our data (Table 2). SNPs rs12520799 in *C5orf20* (*DCNP1*) ( $P=0.038$ ; OR AT=1.10, 95% CI 0.95–1.29; OR TT=1.21, 95% CI 1.01–1.47) and rs16139 in *NPY* ( $P=0.034$ ; OR C allele=0.73, 95% CI 0.55–0.97) replicated the reported effect in the literature. In addition, three SNPs in the *ACE* gene selected from the literature, which were in strong LD with each other ( $r^2=0.78$ – $1.00$ ), were also significant in the GAIN MDD sample

(rs4333:  $P=0.029$ ; rs4329:  $P=0.030$ ; rs4461142:  $P=0.036$ ). However these three SNPs showed effects in the opposite direction compared with previous results<sup>19</sup> (TC Baghai, personal communication). Hence, these SNPs do not represent a true replication.

### Candidate genes from the literature in the GAIN MDD study

As in different populations different SNPs might have a role, we also studied all SNPs present on the genotyping array or available through imputation in an area from 5-kb upstream to 5-kb downstream of each selected candidate gene to cover the promoter and 3' untranslated region, respectively. Of the 57 genes, 49 were covered by one or more SNPs that were present on the Perlegen chip. For another six genes no genotyped SNPs were available but imputed ones were. Neither genotyped nor imputed SNPs were available for two genes (*AVPR1B* and *CHRFAM7A*). In total, 4870 SNPs, of which 820 were genotyped and 4019 were imputed, covered the 55 candidate genes ranging from 1 SNP for *AR* to 642 SNPs for *PDE11A* (Figure 2). We noted a significant correlation of 0.75 between gene size and the number of genotyped SNPs (including genes on the X chromosome) and a correlation of 0.96 between gene size and total number of SNPs (excluding genes on the X chromosome, as SNPs on the X chromosome were not imputed).

For 28 of the 55 genes, one or more SNPs were found to be different between MDD cases and healthy controls in the GAIN MDD GWA scan at a significance level of  $P<0.05$  (Table 3). The remaining 27 genes were not associated with MDD as none of the SNPs reached  $P<0.05$ . The smallest  $P$ -value was observed for SNP rs769178 in the *TNF* gene region ( $P=0.00029$ ; OR T allele = 1.35, 95% CI 1.13–1.63). The minor allele T was observed in 8.0% of the MDD cases and 6.1% of the controls.

With genes covered by a large number of SNP markers, the expected minimal  $P$ -value will decrease purely as a result of chance alone as a function of the number of SNPs. Thus, we used gene-wide  $P$ -values from the permutation procedure that corrects for the number of SNPs within a candidate gene and only SNP rs769178 in *TNF* remained significant ( $P=0.0034$ ; Figure 3 and Table 3). The second strongest associated SNP, which was observed in *DISC1* (rs7533169;  $P$ -value=0.0025), became non-significant after this correction (gene-wide  $P$ -value=0.28), because in this gene there were 491 SNPs and apparently the small  $P$ -value was observed purely based on chance. In addition to the gene-wide significance, we also determined overall significance per SNP accounting for testing 4870 SNPs in 55 candidate genes. In that case none of the SNPs remained significant, not even the *TNF* SNP rs769178 (overall  $P=0.33$ ).

The significance levels per SNP are one way of testing the true value of the selected candidate genes. If many SNPs in a candidate gene are associated, this

**Table 1** Candidate genes and associated polymorphisms or haplotypes based on the systematic literature search

Gene	HUGO name	Chromosome position	Gene size (kb)	Variant	rs-id	Refs			controls			P genot.	OR allele	P genot.	OR genot.	P multi-marker	OR
						N cases	N	P allele	N	P allele	OR allele						
MTHFR	MTHFR	1p36.3	20.3	677C/T	rs1801133	34	32	419	0.005	2.8	0.03	1.90					
5-HT1A	HTR1A	5q11.2-q13	1.3	-1019C/G	rs6295	35	100	89			0.0017	het:1.54 hom:3.75 <sup>a</sup>					
5-HT2A	HTR2A	13q14-q21	62.7	-1438A/G	rs6311	36	129	134	0.0006	1.84 <sup>a</sup>	0.039	0.47					
5-HT2C	HTR2C	Xq24	326.1	102T/C 68G/C = Cys23Ser	rs6313 rs6318	37	71	157	0.018	0.62	0.005	2.4					
5-HT3B	HTR3B	11q23.1	41.7	Tyr129Ser (3B06) Ala154Ala (3B08) IVS6 + 72A > G (3B10) IVS6 + 161C > G (3B11) 3B04 + 3B06 + 3B08 + 3B09 + 3B10 + 3B11 + 3B12	rs1176744 rs2276305 rs2276307 rs2276308 rs1176746 + rs1176744 + rs2276305 + rs3782025	38	189	148	0.007	1.52	0.027	n.a.					
ACE	ACE	17q23.3	20.5	dbSNP4295	rs4295	39	120	131	0.03		0.011	n.a.					
					rs4291	40	513	901	0.006	n.a.	0.023	n.a.					
					rs4305	41	47	90	0.015	2.27	0.031	n.a.				0.002	2.11
					rs4311	42	92	92	<0.0001		0.0011	rec:1.87 <sup>a</sup>				0.0001	n.a.
					rs4329	43	198	229	0.003	1.5 (F) 1.9 (M)	0.001						
					rs4333	44	92	92	0.03	1.59							
					rs4461142	45	642	608			<0.05	n.a.					
					rs867640	46	642	608			<0.05	n.a.					
					C/T in intron 1	47	642	608			<0.01	n.a.					
						48	642	608			<0.05	n.a.					
						49	642	608			= 0.05	n.a.					
						50	642	608			<0.05	n.a.					
						51	642	608			<0.05	n.a.					
						52	642	608			<0.05	n.a.					
						53	198	229	0.003	1.5 (F) 1.9 (M)	0.036	het:0.66 hom:1.61 <sup>a</sup>					
					NA	54	71	83	0.003	0.41 <sup>a</sup>							
					rs5186	55	132	132			0.036	het:0.66 hom:1.61 <sup>a</sup>					
					rs429358 + rs7412	56	42	73	0.04	0.36 <sup>a</sup>							
						57	273	429			<0.0005	het:0.17 hom:0.03 <sup>a</sup>					
					rs33976516	58	89	88	0.026	n.a.	0.014	n.a.					
					rs33933482	59	89	89	0.043	n.a.							
					rs33976516 + rs33933482	60	89	88			0.002	0.52					
						61	89	89			0.0003	0.28					

Table 1 Continued

Gene	HUGO name	Chromosome position	Gene size (kb)	Variant	rs-id	Refs	N cases	N controls	P allele	OR allele	P genot.	OR genot.	P multi-marker	OR			
BDNF	BDNF	11p13	66.9	Val66Met	rs6265	49	110	171	0.001		0.003	2.49					
BCR	BCR	22q11.23	137.7	(GT)n (microsatellite)	NA	50	284	331	0.005	1.7							
					rs2267012	51	465	1097	0.028	n.a.							
					rs3761418	52	329	351	0.0097	1.33							
					rs2267013	52	329	351	0.012	1.32							
					rs2267015	52	329	351	0.044	1.27							
					rs2156921	52	329	351	0.023	1.30							
COMT	COMT	22q11.21	27.2	Val158Met	rs2213172	52	329	351	0.031	1.29							
					rs4680	53	120	628	0.009	1.44							
					rs2075507	54	30	467	0.046	1.79			0.0378	2.12			
					rs2075507 + rs737865 + Val158Met + rs165599	54	30	467									
MCP1	CCL2	17q11.2-q12	1.9	-2518G/A	rs1024611	55	90	114	0.004	1.83	0.027	1.87					
					rs number unknown	16	177	160	0.006	2.58*							
CCKAR	CCKAR	4p15.1-p15.2	9.0	246G/A	rs1799723-rs?-rs1800856-rs1800857	16	177	160					0.002	6.68*			
					rs8191992	56	126	304			0.001	het:0.44 hom:0.47*					
CHRM2	CHRM2	7q31-q35	148.4	1890A/T													
a7nAChR	CHRFAM7A	15q13.1	32.4	2 bp deletion	rs10581632	57	72	103	0.037	n.a.	0.027						
					rs7728378	58	89	88	0.042		0.031						
CRH-BP	CRHBP	5q11.2-q13.3	16.6	CRH-BPs02 CRH-BPs11 CRH-BPs17 + s02 + s14 + s11	rs1875999	58	89	88	0.048		0.043						
					rs3811939 + rs7728378	58	89	88				0.00014	2.21				
CRHR1	CRHR1	17q12-q22	51.5	rs242939	rs242939	59	206	195	0.0008	2.07*	0.0002	het:2.45 hom:1.18*					
CRHR2	CRHR2	7p15.1	29.3	1047G/A	rs2240403	60	89	89	0.04	2.14*	0.03	het:1.21 hom:2.06*					
CYP2C9	CYP2C9	10q24	50.7	CYP2C9*3	rs1057910	61	70	142	<0.05	2.03							
DCNP1	C5orf20	5q31.1	3.1	rs12520799	rs12520799	23	431	433	0.003	2.01*							
					rs number unknown	16	177	160	0.03	0.53*							
DISC1	DISC1	1q42.1	414.5	Ser704Cys (SNP12) SNP1	rs821616 rs6541281	63 63	373 373	717 717	0.005 0.048	1.46 0.75							
DRD1	DRD1	5q35.1	3.5	-2102C/A	rs10063995	16	177	160	0.008	3.69*							
DRD2	DRD2	11q23	65.6	-7054C/A	rs number unknown	16	177	160	0.03	0.53*							
DRD3	DRD3	3q13.3	50.3	Bal I	rs6280	64	36	38	0.014	2.32*	0.027	3.00*					

Table 1 Continued

Gene	HUGO name	Chromosome position	Gene size (kb)	Variant	rs-id	Refs	N cases	N controls	P allele	OR allele	P genot.	OR genot.	P multi-marker	OR
DRD4	DRD4	11p15.5	3.4	48 bp polymorphism	NA	31	49	100	0.011 (4) 0.041 (5)	0.43 <sup>a</sup> 2.79 <sup>a</sup>	0.007 (4/4) 0.035 (4/5)	0.36 <sup>a</sup> 2.99 <sup>a</sup>		
ER-alpha	ESR1	6q25.1	295.7	PvuII	rs2234693	65	126	471	0.004	1.76 <sup>a</sup>	0.004	2.09		
ER-beta	ESR2	14q23.2	111.5	XbaI microsatellite	rs9340799 NA	66	89	126	0.004	1.76 <sup>a</sup>	0.01	het:1.63 hom:3.83 <sup>a</sup>		
GPR50	GPR50	Xq28	4.9	Val606Ile Δ502-505	rs13440581 in complete LD with rs561077	67	102	150	<0.005	0.30 <sup>a</sup>	<0.0005	het:0.40 hom:0.14 <sup>a</sup>		
GABA3	GABRA3	Xq28	284.2	microsatellite	NA	68	136	260	0.0064	1.51 <sup>a</sup>	0.001	2.21		
GMIP	GMIP	19p11-p12	14.2	-525G/A (SNP1) rs2043293 (SNP2) rs2304129 (SNP3) rs880090 (SNP4)	rs3794996 rs2043293 rs2304129 rs880090	69	128	81	<0.0001	0.14	0.0086	1.50		
Gbeta3	GNB3	12p13	7.2	825C/T	rs5443	70	59	59	0.015	1.46	0.001	2.21		
HP	HP	16q22.1	6.4	genotyping based on phenotypes	NA	71	184	158	0.012	2.14	0.008	n.a.		
cPLA2-α	PLA2G4A	1q25	160.1	BanI	rs10798059	72	106	133	0.004	1.67	0.01	rec:0.45 <sup>a</sup>		
CD-MPR	M6PR	12p13	9.3	C/T in 3'UTR of exon 7	rs3318	73	78	111	0.004	1.67	0.008	n.a.		
MAOA	MAOA	Xp11.3	90.7	uVNTR1 (microsatellite)	NA	74	73	68	0.014	1.96	0.029	dom:0.47 <sup>a</sup>		
p75NTR	NGFR	17q21-q22	19.7	EcoRV2 uVNTR1 + EcoRV2	rs1137070 NA	75	228	213	<0.001	2.39	0.008	n.a.		
NPY	NPY	7p15.1	7.7	S205L -399T/C	rs2072446 rs16147	76	133	110	0.009	2.39	0.008	n.a.		
GR	NR3C1	5q31.3	157.6	1128T/C (Leu7Pro) BclI ER22/23EK R23K	rs16139 rs41423247 rs6189 + rs6190 rs6190	77	95	103	0.015	1.97	0.008	n.a.		
PAM	PAM	5q14-q21	163.9	-3211T/C (NR3C1-1) -3211T/C + R23K	rs10482605 rs10482605 + rs6190	78	78	104	0.01	2.3	0.03	1.8	0.008	2.5
PDE10A	PDE10A	6q26	331.1	dbSNP13340364 G>A rs717602 rs220818 rs676389	rs13340364 rs717602 rs220818 rs676389	79	164	164	<0.05	0.54	0.02	n.a.		
						80	48	140	0.03	1.69 <sup>a</sup>	0.02	n.a.		
						81	180	173	0.02	n.a.	0.02	n.a.		
						82	284	331	0.01	n.a.	0.02	n.a.	0.01	4.58
						82	284	331	0.03	n.a.	0.03	n.a.	0.02	n.a.



**Table 1** Continued

Gene	HUGO name	Chromosome position	Gene size (kb)	Variant	rs-id	Refs	N cases	N controls	P allele	OR allele	P genot.	OR genot.	P multi-marker	OR
PDE11A	PDE11A	2q31.2	485.1	rs3770018	rs3770018	82	284	331	0.0005	2.1				
PDE2A	PDE2A	11q13.4	98.2	rs370013	rs370013	82	284	331	0.01					
PDE5A	PDE5A	4q25-q27	134.4	rs3775845	rs3775845	82	284	331	0.007					
PDE6C	PDE6C	10q24	53.1	rs650058	rs650058	82	284	331	0.01					
				rs701865	rs701865	82	284	331	0.03					
PDE9A	PDE9A	21q22.3	121.8	rs729861	rs729861	82	284	331	0.0006	0.6				
P2RX7	P2RX7	12q24	53.2	rs2230912	rs2230912	22	1000	1029	0.0019	1.30				
						25	145	164		1.40	0.02	het:0.99 hom:0.33 <sup>a</sup>		
NET	SLC6A2	16q12.2	47.1	-182T/C	rs2242446							1.86		
SERT	SLC6A4	17q11.1-q12	37.5	5-HTTLPR (44 bp del/ins)	NA	83	112	136			0.019			
						61	70	142	<0.05	2.03				
						71	184	158	0.01	1.82				
						84	68	68	0.021	1.81	0.025	n.a.		
						85	466	836	0.007	1.26	0.015	1.5		
						86	262	475	0.05	1.24				
						87	184	360	0.01	1.40 <sup>a</sup>	0.009	1.39 <sup>a</sup>		
						88	95	156			0.04	2.6		
						89	53	103			0.008	het:2.75 hom:0.87 <sup>a</sup>		
						90	39	193	<0.02	6.51 <sup>a</sup>	<0.004	6.95 <sup>a</sup>		
						91	33	362	0.011	3.73 <sup>a</sup>				
						92	71	99			<0.05	rec:2.50 <sup>a</sup>		
						93	114	120	0.04	0.68				
						94	119	346	0.005	4.44	0.001	n.a.		
						95	74	84					0.0069	2.53
NK1R	TACR1	2p13.1	150.0	5-HTTLPR + VNTR STin2	NA									
LBP-1C	TFCP2	12q13	78.0	dbSNP13013430 C>T	rs13013430	42	92	92	0.04	0.64	0.019	het:2.43 hom:1.39 <sup>a</sup>		
TPH	TPH1	11p15.3-p14	19.8	2236G/A	rs13463	96	180	225	0.016	0.43		het:0.59 hom:1.01 <sup>a</sup>		
				IVS7 + 218A/C	rs1800532	97	91	139			0.032	het:0.78 hom:1.54 <sup>a</sup>		
				SNP1	rs4537731	98	228	253			0.019	het:1.39 hom:2.50 <sup>a</sup>		
				SNP2	rs684302	98	228	253			0.0119	het:1.39 hom:2.50 <sup>a</sup>		
				SNP5	rs1799913	98	228	253	0.0013	1.52 <sup>a</sup>	0.0035	het:1.39 hom:2.50 <sup>a</sup>		
				SNP1 + SNP2 + SNP3 + SNP4 + SNP5 + SNP6	rs4537731 + rs684302 + rs211105 + rs1800532 + rs1799913 + rs7933505	98	228	253			<0.00001	2.11 <sup>a</sup>		

Table 1 Continued

Gene	HUGO name	Chromosome position	Gene size (kb)	Variant	rs-id	Refs	N cases	N controls	P allele	OR allele	P genot.	OR genot.	P multi-marker	OR		
TPH2	TPH2	12q21.1	93.6	SNPB	rs1843809	<sup>99</sup>	300	265	0.0496	1.38						
				SNPE	rs1386494	<sup>99</sup>	300	265	0.0012	0.6						
				SNPB + SNPE + SNPD	rs1843809 + rs1386494 + + rs1386495	<sup>99</sup>	300	265							<0.0001	n.a
				SNP4 = C_245410	rs10784941	<sup>100</sup>	135	315	0.01	0.62 <sup>a</sup>						
TNF-alpha	TNF	6p21.3	2.8	SNP7 = C_15836061	rs2171363	<sup>100</sup>	135	315	0.001	0.51 <sup>a</sup>						
				SNP12 = C_8376042	rs1386486	<sup>100</sup>	135	315	0.01	0.67 <sup>a</sup>						
				1463G/A	unconfirmed	<sup>17</sup>	87	219	<0.001	8.31 <sup>a</sup>						
				variant	variant											
WFS1	WFS1	4p16	33.4	308G/A	rs1800629	<sup>26</sup>	108	125	0.0125	2.24	0.024					
				684C/G = R228R	rs7672995	<sup>101</sup>	177	160	0.007	1.56 <sup>a</sup>						
				1023C/T = F341F	rs56072215	<sup>101</sup>	177	160	0.02	0.55 <sup>a</sup>						
				1185C/T = V395V	rs1801206	<sup>101</sup>	177	160	0.01	1.56 <sup>a</sup>						
				2206G/A = G736S	rs number	<sup>101</sup>	177	160	0.04	0.57 <sup>a</sup>						
				2565G/A = S855S	unknown											
					rs1046316	<sup>101</sup>	177	160	0.04	0.71 <sup>a</sup>						

Abbreviations: dom, dominant; F, female; GAIN, Genetic Association Information Network; GWAS, genome-wide association study; het, heterozygotes; hom, homozygotes; M, male; MDD, major depressive disorder; OR, odds ratio; rec, recessive; rs-id, reference SNP identification number; SNP, single-nucleotide polymorphism.

Sample sizes (N cases/N controls) as well as *P*-values for allelic (*P* allele), genotypic (*P* genotype) and/or multimarker tests (*P* multimarker) are shown. ORs are given whenever possible.

<sup>a</sup>OR was not reported in corresponding article, but could be estimated from genotype counts or frequencies.

<sup>b</sup>SNP is located more than 5 kb away from gene and is not included in our analyses.

<sup>c</sup>Allele was not observed in cases, which resulted in OR=0.00.

**Table 2** Significance of candidate SNPs for MDD identified from the literature within the GAIN MDD GWAS

<i>Gene</i>	<i>SNP</i>	<i>P-value</i>	<i>Genotyped or imputed</i>
<i>ACE</i>	rs4295	0.36	Genotyped
<i>ACE</i>	rs4305	0.11	Imputed
<i>ACE</i>	rs4309	0.057	Imputed
<i>ACE</i>	rs4311	0.21	Imputed
<i>ACE</i>	rs4329	<b>0.030</b>	Imputed
<i>ACE</i>	rs4333	<b>0.029</b>	Imputed
<i>ACE</i>	rs4461142	<b>0.036</b>	Imputed
<i>ACE</i>	rs867640	0.39	Imputed
<i>BCR</i>	rs2156921	0.76	Imputed
<i>BCR</i>	rs2213172 <sup>a</sup>	0.73	Genotyped
<i>BCR</i>	rs2267012 <sup>b</sup>	0.92	Imputed
<i>BCR</i>	rs2267013	0.68	Imputed
<i>BCR</i>	rs2267015	0.89	Imputed
<i>BCR</i>	rs3761418	0.96	Imputed
<i>BDNF</i>	rs6265	0.22	Genotyped
<i>C5orf20</i>	rs12520799	<b>0.038</b>	Genotyped
<i>CCL2</i>	rs1024611	0.50	Imputed
<i>CHRM2</i>	rs8191992	0.13	Imputed
<i>COMT</i>	rs4680	0.55	Genotyped
<i>CRHBP</i>	rs1875999	0.68	Imputed
<i>CRHBP</i>	rs7728378	0.40	Imputed
<i>CRHR1</i>	rs242939	0.22	Imputed
<i>CRHR2</i>	rs2240403	0.86	Genotyped
<i>CYP2C9</i>	rs1057910	0.46	Genotyped
<i>DISC1</i>	rs6541281	0.93	Imputed
<i>DISC1</i>	rs821616	0.58	Genotyped
<i>DRD3</i>	rs6280	0.87	Genotyped
<i>ESR1</i>	rs2234693	0.75	Imputed
<i>ESR1</i>	rs9340799	0.82	Imputed
<i>GMIP</i>	rs2043293	1.00	Imputed
<i>GMIP</i>	rs2304129	0.46	Imputed
<i>GMIP</i>	rs3794996	0.23	Imputed
<i>GMIP</i>	rs880090	0.58	Imputed
<i>GNB3</i>	rs5443 <sup>c</sup>	0.70	Imputed
<i>HTR2A</i>	rs6311	0.57	Imputed
<i>HTR2A</i>	rs6313	0.52	Genotyped
<i>HTR2C</i>	rs6318	0.092	Genotyped
<i>HTR3B</i>	rs1176744	0.50	Genotyped
<i>HTR3B</i>	rs2276307	0.62	Imputed
<i>HTR3B</i>	rs2276308 <sup>d</sup>	0.65	Imputed
<i>MAOA</i>	rs1137070	0.72	Genotyped
<i>MTHFR</i>	rs1801133	0.074	Genotyped
<i>NGFR</i>	rs2072446	0.52	Imputed
<i>NPY</i>	rs16139	<b>0.034</b>	Genotyped
<i>NPY</i>	rs16147	0.82	Imputed
<i>P2RX7</i>	rs2230912	0.42	Genotyped
<i>PAM</i>	rs13340364	0.78	Imputed
<i>PDE10A</i>	rs220818	0.60	Imputed
<i>PDE10A</i>	rs676389	0.48	Imputed
<i>PDE10A</i>	rs717602	0.78	Imputed
<i>PDE11A</i>	rs3770018	0.58	Imputed
<i>PDE5A</i>	rs3775845	0.41	Imputed
<i>PDE6C</i>	rs650058	0.36	Genotyped
<i>PDE6C</i>	rs701865	0.84	Genotyped
<i>SLC6A2</i>	rs2242446	0.37	Imputed
<i>TACR1</i>	rs13013430	0.056	Genotyped
<i>TNF</i>	rs1800629	0.29	Imputed
<i>TPH1</i>	rs1800532	0.066	Imputed
<i>TPH1</i>	rs4537731	0.11	Imputed
<i>TPH1</i>	rs684302	0.071	Imputed

**Table 2** Continued

<i>Gene</i>	<i>SNP</i>	<i>P-value</i>	<i>Genotyped or imputed</i>
<i>TPH2</i>	rs1386486	0.26	Imputed
<i>TPH2</i>	rs1386494	0.61	Imputed
<i>TPH2</i>	rs1843809	0.69	Imputed
<i>TPH2</i>	rs2171363	0.58	Imputed
<i>WFS1</i>	rs1801206	0.18	Imputed

Abbreviations: GAIN, Genetic Association Information Network; GWAS, genome-wide association study; MDD, major depressive disorder; SNP, single-nucleotide polymorphism.

<sup>a</sup>Tagged by rs877590 ( $r^2 = 0.97$ ).

<sup>b</sup>Tagged by rs2267010 ( $r^2 = 1$ ).


<sup>c</sup>Tagged by rs2301339 ( $r^2 = 1$ ).

<sup>d</sup>Tagged by rs12270070 ( $r^2 = 1$ ).

could also indicate potential involvement of a gene in the disorder under study.<sup>20</sup> We noticed, for instance, that in the norepinephrine transporter *SLC6A2* (*NET*), 19 of the 100 SNPs were significant at a *P* level of <0.05 (Table 3). The permutation procedure that tested whether the number of SNPs that were significant at 0.05 was larger than expected, revealed that this finding for *SLC6A2* was indeed significant ( $P = 0.039$ ). Figure 4 shows that the 19 SNPs are scattered in the right half of the gene; 17 of them lie in three independent haplotype blocks and the other 2 SNPs are not in LD with the haplotype blocks or with each other. For none of the other candidate genes this permutation test revealed a significant result.

## Discussion

Several genome-wide linkage studies of MDD have been published (reviewed in Boomsma *et al.*<sup>2</sup>), but the GAIN MDD study is among the first GWAS in MDD.<sup>4,21</sup> We used these GWAS data as a large-scale replication of previously reported significant findings from candidate gene studies in MDD. To this end, we first conducted a systematic review of the literature and identified a total of 57 genes for which a significant association with MDD has been reported at least once. Fifty-five of these genes could be included in our replication study, with either genotyped or imputed SNPs available from the GWAS data. With a sample size (1738 cases and 1802 controls) by far exceeding all previous candidate gene studies (with a mean sample size of 164 MDD cases and 252 controls, and only 1 study with >1000 MDD cases and controls),<sup>22</sup> we found minimal support for involvement in MDD for only 4 out of 55 candidate genes: *C5orf20* ( $P = 0.038$ ), *NPY* ( $P = 0.034$ ), *TNF* ( $P = 0.0034$ ) and *SLC6A2* ( $P = 0.039$ ). Replication of these genes was based on three different statistical approaches. First, the involvement of *C5orf20* (rs12520799) and *NPY* (rs16139) constituted a direct replication at SNP level of previously identified SNPs associated with MDD. Second, studying the selected candidate genes for all SNPs present on



**Figure 2** Coverage by genotyped (black bar) and imputed (grey bar) SNPs and size (white circles) of the 57 selected candidate genes (above: genes with  $\leq 50$  SNPs; below: genes with  $> 50$  SNPs).

the genotyping array, *TNF* (rs76917) was identified as the only gene for which the association with MDD remained gene-wide significant. Third, the potential involvement of *SLC6A2* (*NET*) in MDD was derived on the basis of a statistically significant number of SNPs (that is, 19 out of 100) associated with MDD within this gene, which could not be explained solely by high LD between the SNPs. Note that the previous evidence in the selected studies for the involvement of *C5orf20*, *NPY*, *TNF* and *SLC6A2* in MDD did not stand out as particularly strong; that is, ORs were mostly in the moderate range (minimum OR=0.33 and maximum OR=2.41), and none of the *P*-values in these studies was smaller than  $P < 0.001$ , not even for the largest study on *C5orf20* with a total sample size of 864. *NPY* was the one exception in which the allele was present in 6.3% of the controls but not in the patients, thus suggesting a strong effect (OR=0). However, this was based on only 51 patients.<sup>23–26</sup>

An important question is why so few candidate genes were replicated by our GWAS, whereas the sample size of our study was so much larger than any of the 78 selected candidate gene studies. One possible—and in our opinion most likely—explanation is publication bias; positive results have a better chance of being published than negative results. This would also imply that many previously reported positive findings were actually false-positive findings (type 1 errors), probably amplified by insufficient correction for multiple testing. Testing this, for example, with funnel plots, is not appropriate for the approach followed here as our literature search

strategy did not include negative candidate gene studies. Second, given that MDD is a rather heterogeneous disorder, and that it was diagnosed with different instruments across studies, previous and current study samples may have differed phenotypically (see Hek *et al.*<sup>27</sup> for a discussion of this point). A third explanation is that associations between genes and etiologically complex diseases may depend on genetic (gene  $\times$  gene interactions or epistasis) and environmental (gene  $\times$  environment interactions) contexts, which may differ in samples from different populations.<sup>28</sup> Thus, previous and current samples may have been different genetically or in their contextual factors.

Given the modest support for the replicated candidate genes (one *P*-value  $< 0.01$  and three *P*-values  $< 0.05$ ) it is possible that even the four replications of our analysis are false-positive results. With the number of SNPs ( $n=65$ ) been tested and under the null hypothesis of no true associations of any of the candidate SNPs in previous studies, one would expect three significant findings. Thus, it is possible that our two significant findings in *C5orf20* and *NPY* are false-positive results. In addition, under a similar null hypothesis for the 55 candidate genes, two significant findings were expected, implying that the single gene-wide significant result of *TNF* might also be a false-positive result. This is supported by the lack of overall significance for the *TNF* SNP ( $P=0.33$ ). And finally, the significant finding for *SLC6A2*, the only gene showing a larger number of significant SNPs than expected, might also be a chance finding.


**Table 3** Gene-wide results for the 57 candidate genes identified by the systematic literature search

Gene	Genotyped	Imputed	Total	Minimal P (analytical)	Minimal P (permuted)	Gene- wide P	Overall P	rs-id	SNPs at P < 0.05	P-value SNPs
TNF	2	18	20	<b>0.00029</b>	<b>0.00094</b>	<b>0.0034</b>	0.34	rs769178	1	0.32
DISC1	105	386	491	<b>0.0025</b>	<b>0.0015</b>	0.29	0.95	rs7533169	18	0.56
SLC6A2	25	75	100	<b>0.0061</b>	<b>0.0058</b>	0.19	1.00	rs5564	19	<b>0.039</b>
DRD2	12	68	80	<b>0.0079</b>	<b>0.0077</b>	0.15	1.00	rs7131056	3	0.32
HTR3B	6	33	39	<b>0.0081</b>	<b>0.0077</b>	0.11	1.00	rs7945926	3	0.21
CHRM2	43	157	200	<b>0.0097</b>	<b>0.0081</b>	0.32	1.00	rs6944132	3	0.54
MTHFR	6	37	43	<b>0.0098</b>	<b>0.0080</b>	0.11	1.00	rs3737967	4	0.18
PDE10A	107	392	499	<b>0.013</b>	<b>0.014</b>	0.74	1.00	rs7741623	8	0.80
APOE	0	5	5	<b>0.015</b>	<b>0.012</b>	0.057	1.00	rs10119	1	0.11
TACR1	31	141	172	<b>0.016</b>	<b>0.019</b>	0.50	1.00	rs1012686	24	0.10
TPH2	21	134	155	<b>0.017</b>	<b>0.016</b>	0.29	1.00	rs17722134	6	0.29
PDE11A	66	576	642	<b>0.018</b>	<b>0.016</b>	0.63	1.00	rs6433706	11	0.57
DRD3	13	41	54	<b>0.018</b>	<b>0.021</b>	0.23	1.00	rs1800827	3	0.27
PDE5A	7	108	115	<b>0.019</b>	<b>0.020</b>	0.20	1.00	rs13104219	21	0.08
NR3C1	14	60	74	<b>0.024</b>	<b>0.025</b>	0.37	1.00	rs4607376	12	0.11
MAOA	5	0	5	<b>0.026</b>	<b>0.028</b>	0.078	1.00	rs5906883	1	0.11
ESR1	50	272	322	<b>0.027</b>	<b>0.025</b>	0.84	1.00	rs17081749	1	0.93
ACE	3	28	31	<b>0.028</b>	<b>0.031</b>	0.21	1.00	rs4344	8	0.08
NPY	6	38	44	<b>0.030</b>	<b>0.031</b>	0.29	1.00	rs1073317	6	0.15
PDE9A	25	133	158	<b>0.033</b>	<b>0.031</b>	0.69	1.00	rs13046735	1	0.75
HTR2A	35	100	135	<b>0.037</b>	<b>0.037</b>	0.72	1.00	rs2770292	3	0.56
C5orf20	10	35	45	<b>0.038</b>	<b>0.038</b>	0.36	1.00	rs12520799	5	0.13
TPH1	6	24	30	<b>0.040</b>	<b>0.038</b>	0.25	1.00	rs10741734	2	0.19
PLA2G4A	26	92	118	<b>0.040</b>	<b>0.042</b>	0.68	1.00	rs7555140	1	0.68
BCR	14	110	124	<b>0.041</b>	<b>0.045</b>	0.69	1.00	rs11090231	1	0.67
ESR2	43	155	198	<b>0.042</b>	<b>0.040</b>	0.80	1.00	rs1255949	2	0.68
P2RX7	21	87	108	<b>0.047</b>	<b>0.049</b>	0.62	1.00	rs11065464	1	0.57
AGTR1	8	44	52	0.054	<b>0.049</b>	0.58	1.00	rs12721331	0	0.77
PDE6C	18	52	70	0.055	0.056	0.71	1.00	rs10882288	0	0.84
CYP2C9	8	47	55	0.057	0.056	0.48	1.00	rs1799853	0	0.72
CCKAR	4	17	21	0.060	0.058	0.33	1.00	rs11726159	0	0.64
NGFR	4	16	20	0.065	0.064	0.55	1.00	rs741073	0	0.73
TFCP2	4	59	63	0.066	0.064	0.53	1.00	rs2029686	0	0.72
CRHR2	12	35	47	0.083	0.085	0.70	1.00	rs2267717	0	0.77
GPR50	3	0	3	0.090	0.092	0.24	1.00	rs5969987	0	0.57
HTR2C	10	0	10	0.092	0.091	0.39	1.00	rs6318	0	0.62
CRHBP	3	9	12	0.093	0.092	0.43	1.00	rs7721799	0	0.63
PAM	7	92	99	0.096	0.090	0.81	1.00	rs1650500	0	0.79
SLC6A4	6	26	32	0.11	0.11	0.72	1.00	rs12150465	0	0.72
M6PR	3	21	24	0.11	0.10	0.54	1.00	rs1805729	0	0.65
BDNF	7	29	36	0.11	0.11	0.60	1.00	rs11030104	0	0.68
GABRA3	10	0	10	0.11	0.10	0.50	1.00	rs994423	0	0.63
WFS1	3	55	58	0.12	0.13	0.51	1.00	rs4689393	0	0.63
COMT	5	31	36	0.13	0.13	0.78	1.00	rs2871047	0	0.72
CCL2	0	9	9	0.14	0.14	0.55	1.00	rs2857655	0	0.62
DRD1	5	16	21	0.17	0.17	0.79	1.00	rs5326	0	0.69
PDE2A	16	62	78	0.20	0.19	0.97	1.00	rs11604811	0	0.81
CRHR1	4	42	46	0.22	0.22	0.86	1.00	rs242939	0	0.69
AR	1	0	1	0.23	0.23	0.23	1.00	rs2207040	0	0.52
GMP	0	10	10	0.23	0.23	0.83	1.00	rs12003	0	0.66
DRD4	0	6	6	0.29	0.30	0.74	1.00	rs3758653	0	0.59
ACSL4	2	0	2	0.31	0.30	0.50	1.00	rs7887981	0	0.55
HP	1	5	6	0.32	0.32	0.77	1.00	rs5475	0	0.59
GNB3	3	10	13	0.39	0.39	0.96	1.00	rs3759348	0	0.23
HTR1A	0	5	5	0.54	0.55	0.79	1.00	rs10042486	0	0.55
AVPR1B	0	0	0	NA	NA	NA	NA	NA	NA	NA
CHRFAM7A	0	0	0	NA	NA	NA	NA	NA	NA	NA

Abbreviations: NA, not applicable; SNP, single-nucleotide polymorphism; rs-id, reference SNP identification number. Analytical *P*-values were determined using SNPTTEST for the autosomes and logistic regression in PLINK for the X chromosome. Permutations ( $N=10\,000$ ) were performed to determine uncorrected SNP *P*-values, gene-wide corrected *P*-values, overall corrected *P*-values and *P*-values for number of SNPs with  $P < 0.05$ . All *P*-values were calculated assuming an additive model. *P*-values of  $< 0.05$  are shown in bold.

On the other hand, the above calculations may be too conservative when assuming that at least some of the previously found associations of candidate genes with MDD were true findings. In that case, our approach of replicating candidate genes is more or less hypothesis driven, thus not requiring the same multiple testing penalty as the genome-wide approach.<sup>4</sup> Nonetheless, the few replications in our study are rather sobering and to uncover whether the replicated SNPs and genes are truly associated to MDD, confirmation in independent samples is crucial. As such, meta-analytical results from the Psychiatric GWAS Consortium are also eagerly awaited.<sup>29</sup>


From the most recent meta-analysis of genetic studies on MDD, Lopez-Leon *et al.*<sup>9</sup> concluded that statistically significant evidence exists for six MDD



**Figure 3** Observed minimal  $P$ -value versus the expected minimal  $P$ -value based on the number of SNPs in the gene.

susceptibility genes, that is, *APOE*, *DRD4*, *GNB3*, *MTHFR*, *SLC6A3* and *SLC6A4*. Our study offers little support for these genes. Given our sample size, we had 80% power to detect ORs of  $>1.15$  for allele frequencies  $>5\%$  under an additive disease model. All of the above genes meet these criteria. The association of *GNB3* and *MTHFR* with MDD was directly tested but could not be replicated in the GAIN MDD sample, although the effect of *MTHFR* showed a trend in the expected direction ( $P=0.074$ ; OR TT versus CC = 1.14). In addition, *APOE*, *DRD4* and *SLC6A4* were not associated with MDD in our study, but the previously identified genetic variants were length polymorphisms instead of SNPs. Hence, these could not directly be tested and as it is unsure whether these variants are tagged by the SNPs in the corresponding genes, we cannot refute the associations. *SLC6A3* was not identified in our literature search as a candidate gene for MDD, because individual studies did not report significant effects for this gene<sup>30,31</sup> and it only reached significance in the pooled meta-analysis. We also tested for significance in our sample *post hoc* and did not find any association (59 SNPs, most significant SNP: rs27072,  $P=0.096$ ).

In the context of the non-replication of many of the selected candidate genes as discussed earlier, two limitations of our study need to be noted. First, we did not have direct or indirect information on one-third of the candidate SNPs, as these were not present on the genotyping chip and could not be imputed using the HapMap CEU data. Second, we were unable to test candidate length polymorphisms previously associated with MDD. Among these was the 44-bp insertion/deletion polymorphism (or 5-HTTLPR) in the promoter region of the serotonin transporter gene *SLC6A4* (*SERT*). Length polymorphisms are often difficult to tag with single SNPs because LD information between SNPs and length polymorphisms is either unavailable or LD with SNPs is insufficiently strong.<sup>32</sup> However, a recent study by Wray *et al.*<sup>33</sup> identified a two-SNP haplotype proxy for 5-HTTLPR.



**Figure 4** Association of *SLC6A2* (NET) with MDD. Along the x axis the location of all 100 SNPs within 5 kb of *SLC6A2* is shown and the  $-\log_{10}(P\text{-value})$  is on the y axis. A total of 19 SNPs have a  $P$ -value of  $<0.05$ , which is significantly more than expected ( $P=0.039$ ). Of these, 17 lie in three haplotype blocks (squares–upward triangles (light grey) and downward triangles (dark grey)) ( $r^2$  values HapMap CEU: squares–upward triangles: 0.020; squares–downward triangles: 0.661; and upward triangles–downward triangles: 0.013). The other two (black diamonds) SNPs are not in LD with the haplotype blocks or with each other. The light grey line shows the recombination rate in this area (axis on the right).

In conclusion, the GWAS data of the GAIN MDD study failed to replicate all but four of the previously reported candidate gene associations with MDD. However, given the number of candidate SNPs and genes that were tested, even these significant may well be false-positives, implying that we found no replication at all. This might point to publication bias and false-positive findings in previous studies, and also to heterogeneity of the MDD phenotype as well as variations in contextual genetic or environmental factors.

### Conflict of interest

The authors declare no conflict of interest.

### Acknowledgments

We thank Ting Wu for making the flow diagram in Figure 1.

### References

- 1 Manolio TA, Rodriguez LL, Brooks L, Abecasis G, Ballinger D, Daly M *et al*. New models of collaboration in genome-wide association studies: the Genetic Association Information Network. *Nat Genet* 2007; **39**: 1045–1051.
- 2 Boomsma DI, Willemsen G, Sullivan PF, Heutink P, Meijer P, Sondervan D *et al*. Genome-wide association of major depression: description of samples for the GAIN Major Depressive Disorder Study: NTR and NESDA biobank projects. *Eur J Hum Genet* 2008; **16**: 335–342.
- 3 The Wellcome Trust Case-Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 2007; **447**: 661–678.
- 4 Sullivan PF, de Geus EJ, Willemsen G, James MR, Smit JH, Zandbelt T *et al*. Genome-wide association for major depressive disorder: a possible role for the presynaptic protein piccolo. *Mol Psychiatry* 2009; **14**: 359–375.
- 5 Arking DE, Pfeufer A, Post W, Kao WH, Newton-Cheh C, Ikeda M *et al*. A common genetic variant in the NOS1 regulator NOS1AP modulates cardiac repolarization. *Nat Genet* 2006; **38**: 644–651.
- 6 Roeder K, Bacanu SA, Wasserman L, Devlin B. Using linkage genome scans to improve power of association in genome scans. *Am J Hum Genet* 2006; **78**: 243–252.
- 7 Hirschhorn JN, Lohmueller K, Byrne E, Hirschhorn K. A comprehensive review of genetic association studies. *Genet Med* 2002; **4**: 45–61.
- 8 Munafò MR. Candidate gene studies in the 21st century: meta-analysis, mediation, moderation. *Genes Brain Behav* 2006; **5**(Suppl 1): 3–8.
- 9 Lopez-Leon S, Janssens AC, Gonzalez-Zuloeta Ladd AM, Del-Favero J, Claes SJ, Oostra BA *et al*. Meta-analyses of genetic studies on major depressive disorder. *Mol Psychiatry* 2008; **13**: 772–785.
- 10 Penninx BW, Beekman AT, Smit JH, Zitman FG, Nolen W, Spinhoven P *et al*. The Netherlands Study of Depression and Anxiety (NESDA): rationale, objectives and methods. *Int J Methods Psychiatr Res* 2008; **17**: 121–140.
- 11 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. American Psychiatric Association: Washington, DC, 1994.
- 12 World Health Organization. *Composite International Diagnostic Interview (CIDI), Version 2.1..* World Health Organization: Geneva, Switzerland, 1997.
- 13 Marchini J, Howie B, Myers S, McVean G, Donnelly P. A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat Genet* 2007; **39**: 906–913.
- 14 Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D *et al*. PLINK: a tool set for whole-genome association and

- population-based linkage analyses. *Am J Hum Genet* 2007; **81**: 559–575.
- 15 Maes M, Delanghe J, Scharpe S, Meltzer HY, Cosyns P, Suy E *et al*. Haptoglobin phenotypes and gene frequencies in unipolar major depression. *Am J Psychiatry* 1994; **151**: 112–116.
- 16 Koks S, Nikopoulou T, Koido K, Maron E, Altmae S, Heinaste E *et al*. Analysis of SNP profiles in patients with major depressive disorder. *Int J Neuropsychopharmacol* 2006; **9**: 167–174.
- 17 Zhang X, Gainetdinov RR, Beaulieu JM, Sotnikova TD, Burch LH, Williams RB *et al*. Loss-of-function mutation in tryptophan hydroxylase-2 identified in unipolar major depression. *Neuron* 2005; **45**: 11–16.
- 18 Blakely RD. Overview: a rare opportunity or just one less reason to be depressed. *Neuron* 2005; **48**: 701–702; author reply 705–706.
- 19 Baghai TC, Binder EB, Schule C, Salyakina D, Eser D, Lucae S *et al*. Polymorphisms in the angiotensin-converting enzyme gene are associated with unipolar depression, ACE activity and hypercortisolism. *Mol Psychiatry* 2006; **11**: 1003–1015.
- 20 Arking DE, Khera A, Xing C, Kao WH, Post W, Boerwinkle E *et al*. Multiple independent genetic factors at NOS1AP modulate the QT interval in a multi-ethnic population. *PLoS ONE* 2009; **4**: e4333.
- 21 Muglia P, Tozzi F, Galwey NW, Francks C, Upmanyu R, Kong XQ *et al*. Genome-wide association study of recurrent major depressive disorder in two European case-control cohorts. *Mol Psychiatry*; advance online publication, 23 December 2008; doi:10.1038/mp.2008.131.
- 22 Lucae S, Salyakina D, Barden N, Harvey M, Gagne B, Labbe M *et al*. P2RX7, a gene coding for a purinergic ligand-gated ion channel, is associated with major depressive disorder. *Hum Mol Genet* 2006; **15**: 2438–2445.
- 23 Willis-Owen SA, Shifman S, Copley RR, Flint J. DCNP1: a novel candidate gene for major depression. *Mol Psychiatry* 2006; **11**: 121–122.
- 24 Heilig M, Zachrisson O, Thorsell A, Ehnvall A, Mottagui-Tabar S, Sjogren M *et al*. Decreased cerebrospinal fluid neuropeptide Y (NPY) in patients with treatment refractory unipolar major depression: preliminary evidence for association with preproNPY gene polymorphism. *J Psychiatr Res* 2004; **38**: 113–121.
- 25 Inoue K, Itoh K, Yoshida K, Shimizu T, Suzuki T. Positive association between T-182C polymorphism in the norepinephrine transporter gene and susceptibility to major depressive disorder in a Japanese population. *Neuropsychobiology* 2004; **50**: 301–304.
- 26 Jun TY, Pae CU, Hoon H, Chae JH, Bahk WM, Kim KS *et al*. Possible association between -G308A tumour necrosis factor- $\alpha$  gene polymorphism and major depressive disorder in the Korean population. *Psychiatr Genet* 2003; **13**: 179–181.
- 27 Hek K, Mulder CL, Luijendijk HJ, van Duijn CM, Hofman A, Uitterlinden AG *et al*. The PCLO gene and depressive disorders: replication in a population-based study. *Hum Mol Genet* 2009; **19**: 731–734.
- 28 Martinez FD. Gene-environment interactions in asthma: with apologies to William of Ockham. *Proc Am Thorac Soc* 2007; **4**: 26–31.
- 29 Psychiatric GWAS Consortium Steering Committee. A framework for interpreting genome-wide association studies of psychiatric disorders. *Mol Psychiatry* 2009; **14**: 10–17.
- 30 Frisch A, Postilnick D, Rockah R, Michaelovsky E, Postilnick S, Birman E *et al*. Association of unipolar major depressive disorder with genes of the serotonergic and dopaminergic pathways. *Mol Psychiatry* 1999; **4**: 389–392.
- 31 Manki H, Kanba S, Muramatsu T, Higuchi S, Suzuki E, Matsushita S *et al*. Dopamine D2, D3 and D4 receptor and transporter gene polymorphisms and mood disorders. *J Affect Disord* 1996; **40**: 7–13.
- 32 Su S, Zhao J, Bremner JD, Miller AH, Tang W, Bouzyk M *et al*. Serotonin transporter gene, depressive symptoms and interleukin-6. *Circ Cardiovasc Genet* 2009; **2**: 614–620.
- 33 Wray NR, James MR, Gordon SD, Dumenil T, Ryan L, Coventry WL *et al*. Accurate, large-scale genotyping of 5HTTLPR and flanking single nucleotide polymorphisms in an association study of depression, anxiety, and personality measures. *Biol Psychiatry* 2009; **66**: 468–476.

- 34 Arinami T, Yamada N, Yamakawa-Kobayashi K, Hamaguchi H, Toru M. Methylenetetrahydrofolate reductase variant and schizophrenia/depression. *Am J Med Genet* 1997; **74**: 526–528.
- 35 Kelly CB, McDonnell AP, Johnston TG, Mulholland C, Cooper SJ, McMaster D et al. The MTHFR C677 T polymorphism is associated with depressive episodes in patients from Northern Ireland. *J Psychopharmacol* 2004; **18**: 567–571.
- 36 Lemonde S, Turecki G, Bakish D, Du L, Hrdina PD, Bown CD et al. Impaired repression at a 5-hydroxytryptamine 1A receptor gene polymorphism associated with major depression and suicide. *J Neurosci* 2003; **23**: 8788–8799.
- 37 Choi MJ, Kang RH, Ham BJ, Jeong HY, Lee MS. Serotonin receptor 2A gene polymorphism (-1438A/G) and short-term treatment response to citalopram. *Neuropsychobiology* 2005; **52**: 155–162.
- 38 Choi MJ, Lee HJ, Lee HJ, Ham BJ, Cha JH, Ryu SH et al. Association between major depressive disorder and the -1438A/G polymorphism of the serotonin 2A receptor gene. *Neuropsychobiology* 2004; **49**: 38–41.
- 39 Du L, Bakish D, Lapierre YD, Ravindran AV, Hrdina PD. Association of polymorphism of serotonin 2A receptor gene with suicidal ideation in major depressive disorder. *Am J Med Genet* 2000; **96**: 56–60.
- 40 Lerer B, Macciardi F, Segman RH, Adolfsson R, Blackwood D, Blairy S et al. Variability of 5-HT<sub>2C</sub> receptor cys23ser polymorphism among European populations and vulnerability to affective disorder. *Mol Psychiatry* 2001; **6**: 579–585.
- 41 Yamada K, Hattori E, Iwayama Y, Ohnishi T, Ohba H, Toyota T et al. Distinguishable haplotype blocks in the HTR3A and HTR3B region in the Japanese reveal evidence of association of HTR3B with female major depression. *Biol Psychiatry* 2006; **60**: 192–201.
- 42 Mendlewicz J, Oswald P, Claes S, Massat I, Souery D, Van Broeckhoven C et al. Patient-control association study of substance P-related genes in unipolar and bipolar affective disorders. *Int J Neuropsychopharmacol* 2005; **8**: 505–513.
- 43 Covault J, Pettinati H, Moak D, Mueller T, Kranzler HR. Association of a long-chain fatty acid-CoA ligase 4 gene polymorphism with depression and with enhanced niacin-induced dermal erythema. *Am J Med Genet B Neuropsychiatr Genet* 2004; **127B**: 42–47.
- 44 Su QR, Su LY, Su HR, Chen Q, Ren GY, Yin Y et al. Polymorphisms of androgen receptor gene in childhood and adolescent males with first-onset major depressive disorder and association with related symptomatology. *Int J Neurosci* 2007; **117**: 903–917.
- 45 Saab YB, Gard PR, Yeoman MS, Mfarrej B, El-Moalem H, Ingram MJ. Renin-angiotensin-system gene polymorphisms and depression. *Prog Neuropsychopharmacol Biol Psychiatry* 2007; **31**: 1113–1118.
- 46 Holmes C, Russ C, Kirov G, Aitchison KJ, Powell JF, Collier DA et al. Apolipoprotein E: depressive illness, depressive symptoms, and Alzheimer's disease. *Biol Psychiatry* 1998; **43**: 159–164.
- 47 Fan PL, Chen CD, Kao WT, Shu BC, Lung FW. Protective effect of the apo epsilon2 allele in major depressive disorder in Taiwanese. *Acta Psychiatr Scand* 2006; **113**: 48–53.
- 48 van West D, Del-Favero J, Aulchenko Y, Oswald P, Souery D, Forsgren T et al. A major SNP haplotype of the arginine vasopressin 1B receptor protects against recurrent major depression. *Mol Psychiatry* 2004; **9**: 287–292.
- 49 Hwang JP, Tsai SJ, Hong CJ, Yang CH, Lirng JF, Yang YM. The Val66Met polymorphism of the brain-derived neurotrophic-factor gene is associated with geriatric depression. *Neurobiol Aging* 2006; **27**: 1834–1837.
- 50 Ribeiro L, Busnello JV, Cantor RM, Whelan F, Whittaker P, Deloukas P et al. The brain-derived neurotrophic factor rs6265 (Val66Met) polymorphism and depression in Mexican-Americans. *NeuroReport* 2007; **18**: 1291–1293.
- 51 Schumacher J, Jamra RA, Becker T, Ohlraun S, Klopp N, Binder EB et al. Evidence for a relationship between genetic variants at the brain-derived neurotrophic factor (BDNF) locus and major depression. *Biol Psychiatry* 2005; **58**: 307–314.
- 52 Hashimoto R, Okada T, Kato T, Kosuga A, Tatsumi M, Kamijima K et al. The breakpoint cluster region gene on chromosome 22q11 is associated with bipolar disorder. *Biol Psychiatry* 2005; **57**: 1097–1102.
- 53 Massat I, Souery D, Del-Favero J, Nothen M, Blackwood D, Muir W et al. Association between COMT (Val158Met) functional polymorphism and early onset in patients with major depressive disorder in a European multicenter genetic association study. *Mol Psychiatry* 2005; **10**: 598–605.
- 54 Funke B, Malhotra AK, Finn CT, Plocik AM, Lake SL, Lencz T et al. COMT genetic variation confers risk for psychotic and affective disorders: a case control study. *Behav Brain Funct* 2005; **1**: 19.
- 55 Pae CU, Yu HS, Kim TS, Lee CU, Lee SJ, Jun TY et al. Monocyte chemoattractant protein-1 (MCP1) promoter -2518 polymorphism may confer a susceptibility to major depressive disorder in the Korean population. *Psychiatry Res* 2004; **127**: 279–281.
- 56 Comings DE, Wu S, Rostamkhani M, McGue M, Iacono WG, MacMurray JP. Association of the muscarinic cholinergic 2 receptor (CHRM2) gene with major depression in women. *Am J Med Genet* 2002; **114**: 527–529.
- 57 Lai IC, Hong CJ, Tsai SJ. Association study of nicotinic-receptor variants and major depressive disorder. *J Affect Disord* 2001; **66**: 79–82.
- 58 Claes S, Villafuerte S, Forsgren T, Sluijs S, Del-Favero J, Adolfsson R et al. The corticotropin-releasing hormone binding protein is associated with major depression in a population from Northern Sweden. *Biol Psychiatry* 2003; **54**: 867–872.
- 59 Liu Z, Zhu F, Wang G, Xiao Z, Wang H, Tang J et al. Association of corticotropin-releasing hormone receptor1 gene SNP and haplotype with major depression. *Neurosci Lett* 2006; **404**: 358–362.
- 60 Villafuerte SM, Del-Favero J, Adolfsson R, Souery D, Massat I, Mendlewicz J et al. Gene-based SNP genetic association study of the corticotropin-releasing hormone receptor-2 (CRHR2) in major depression. *Am J Med Genet* 2002; **114**: 222–226.
- 61 Dorado P, Penas-Lledo EM, Gonzalez AP, Caceres MC, Coboleda J, Llerena A. Increased risk for major depression associated with the short allele of the serotonin transporter promoter region (5-HTTLPR-S) and the CYP2C9\*3 allele. *Fundam Clin Pharmacol* 2007; **21**: 451–453.
- 62 Llerena A, Berecz R, Dorado P, Gonzalez AP, Penas LEM, De La Rubia A. CYP2C9 gene and susceptibility to major depressive disorder. *Pharmacogenomics J* 2003; **3**: 300–302.
- 63 Hashimoto R, Numakawa T, Ohnishi T, Kumamaru E, Yagasaki Y, Ishimoto T et al. Impact of the DISC1 Ser704Cys polymorphism on risk for major depression, brain morphology and ERK signaling. *Hum Mol Genet* 2006; **15**: 3024–3033.
- 64 Dikeos DG, Papadimitriou GN, Avramopoulos D, Karadima G, Daskalopoulou EG, Souery D et al. Association between the dopamine D3 receptor gene locus (DRD3) and unipolar affective disorder. *Psychiatr Genet* 1999; **9**: 189–195.
- 65 Serretti A, Lilli R, Di Bella D, Bertelli S, Nobile M, Novelli E et al. Dopamine receptor D4 gene is not associated with major psychoses. *Am J Med Genet* 1999; **88**: 486–491.
- 66 Tsai SJ, Wang YC, Hong CJ, Chiu HJ. Association study of oestrogen receptor alpha gene polymorphism and suicidal behaviours in major depressive disorder. *Psychiatr Genet* 2003; **13**: 19–22.
- 67 Geng YG, Su QR, Su LY, Chen Q, Ren GY, Shen SQ et al. Comparison of the polymorphisms of androgen receptor gene and estrogen alpha and beta gene between adolescent females with first-onset major depressive disorder and controls. *Int J Neurosci* 2007; **117**: 539–547.
- 68 Thomson PA, Wray NR, Thomson AM, Dunbar DR, Grassie MA, Condie A et al. Sex-specific association between bipolar affective disorder in women and GPR50, an X-linked orphan G protein-coupled receptor. *Mol Psychiatry* 2005; **10**: 470–478.
- 69 Henkel V, Baghai TC, Eser D, Zill P, Mergl R, Zwanzger P et al. The gamma amino butyric acid (GABA) receptor alpha-3 subunit gene polymorphism in unipolar depressive disorder: a genetic association study. *Am J Med Genet B Neuropsychiatr Genet* 2004; **126B**: 82–87.
- 70 Tadokoro K, Hashimoto R, Tatsumi M, Kosuga A, Kamijima K, Kunugi H. The Gem interacting protein (GMIP) gene is associated with major depressive disorder. *Neurogenetics* 2005; **6**: 127–133.
- 71 Cao MQ, Hu SY, Zhang CH, Xia DS. Study on the interrelationship between 5-HTTLPR/G-protein beta3 subunit (C825 T) polymorphisms and depressive disorder. *Psychiatr Genet* 2007; **17**: 233–238.



- 72 Lee HJ, Cha JH, Ham BJ, Han CS, Kim YK, Lee SH *et al*. Association between a G-protein beta 3 subunit gene polymorphism and the symptomatology and treatment responses of major depressive disorders. *Pharmacogenomics J* 2004; **4**: 29–33.
- 73 Zill P, Baghai TC, Zwanzger P, Schule C, Minov C, Riedel M *et al*. Evidence for an association between a G-protein beta3-gene variant with depression and response to antidepressant treatment. *NeuroReport* 2000; **11**: 1893–1897.
- 74 Pae CU, Yu HS, Kim JJ, Lee CU, Lee SJ, Lee KU *et al*. BanI polymorphism of the cytosolic phospholipase A2 gene and mood disorders in the Korean population. *Neuropsychobiology* 2004; **49**: 185–188.
- 75 Kolsch H, Ptok U, Majores M, Schmitz S, Rao ML, Maier W *et al*. Putative association of polymorphism in the mannose 6-phosphate receptor gene with major depression and Alzheimer's disease. *Psychiatr Genet* 2004; **14**: 97–100.
- 76 Schulze TG, Muller DJ, Krauss H, Scherk H, Ohlraun S, Sygailo YV *et al*. Association between a functional polymorphism in the monoamine oxidase A gene promoter and major depressive disorder. *Am J Med Genet* 2000; **96**: 801–803.
- 77 Yu YW, Tsai SJ, Hong CJ, Chen TJ, Chen MC, Yang CW. Association study of a monoamine oxidase a gene promoter polymorphism with major depressive disorder and antidepressant response. *Neuropsychopharmacology* 2005; **30**: 1719–1723.
- 78 Du L, Bakish D, Ravindran A, Hrdina PD. MAO-A gene polymorphisms are associated with major depression and sleep disturbance in males. *NeuroReport* 2004; **15**: 2097–2101.
- 79 Kunugi H, Hashimoto R, Yoshida M, Tatsumi M, Kamijima K. A missense polymorphism (S205 L) of the low-affinity neurotrophin receptor p75NTR gene is associated with depressive disorder and attempted suicide. *Am J Med Genet B Neuropsychiatr Genet* 2004; **129B**: 44–46.
- 80 van Rossum EF, Binder EB, Majer M, Koper JW, Ising M, Modell S *et al*. Polymorphisms of the glucocorticoid receptor gene and major depression. *Biol Psychiatry* 2006; **59**: 681–688.
- 81 van West D, Van Den Eede F, Del-Favero J, Souery D, Norrback KF, Van Duijn C *et al*. Glucocorticoid receptor gene-based SNP analysis in patients with recurrent major depression. *Neuropsychopharmacology* 2006; **31**: 620–627.
- 82 Wong ML, Whelan F, Deloukas P, Whittaker P, Delgado M, Cantor RM *et al*. Phosphodiesterase genes are associated with susceptibility to major depression and antidepressant treatment response. *Proc Natl Acad Sci USA* 2006; **103**: 15124–15129.
- 83 Ryu SH, Lee SH, Lee HJ, Cha JH, Ham BJ, Han CS *et al*. Association between norepinephrine transporter gene polymorphism and major depression. *Neuropsychobiology* 2004; **49**: 174–177.
- 84 Nobile M, Cataldo MG, Giorda R, Battaglia M, Baschiroto C, Bellina M *et al*. A case-control and family-based association study of the 5-HTTLPR in pediatric-onset depressive disorders. *Biol Psychiatry* 2004; **56**: 292–295.
- 85 Hoefgen B, Schulze TG, Ohlraun S, von Widdern O, Hofels S, Gross M *et al*. The power of sample size and homogenous sampling: association between the 5-HTTLPR serotonin transporter polymorphism and major depressive disorder. *Biol Psychiatry* 2005; **57**: 247–251.
- 86 Cervilla JA, Rivera M, Molina E, Torres-Gonzalez F, Bellon JA, Moreno B *et al*. The 5-HTTLPR s/s genotype at the serotonin transporter gene (SLC6A4) increases the risk for depression in a large cohort of primary care attendees: the PREDICT-gene study. *Am J Med Genet B Neuropsychiatr Genet* 2006; **141B**: 912–917.
- 87 Grunblatt E, Loffler C, Zehetmayer S, Jungwirth S, Tragl KH, Riederer P *et al*. Association study of the 5-HTTLPR polymorphism and depression in 75-year-old nondemented subjects from the Vienna Transdanube Aging (VITA) study. *J Clin Psychiatry* 2006; **67**: 1373–1378.
- 88 Munafo MR, Clark TG, Roberts KH, Johnstone EC. Neuroticism mediates the association of the serotonin transporter gene with lifetime major depression. *Neuropsychobiology* 2006; **53**: 1–8.
- 89 Mann JJ, Huang YY, Underwood MD, Kassir SA, Oppenheim S, Kelly TM *et al*. A serotonin transporter gene promoter polymorphism (5-HTTLPR) and prefrontal cortical binding in major depression and suicide. *Arch Gen Psychiatry* 2000; **57**: 729–738.
- 90 Ogilvie AD, Battersby S, Bubbs VJ, Fink G, Harmar AJ, Goodwin GM *et al*. Polymorphism in serotonin transporter gene associated with susceptibility to major depression. *Lancet* 1996; **347**: 731–733.
- 91 Liu W, Gu N, Feng G, Li S, Bai S, Zhang J *et al*. Tentative association of the serotonin transporter with schizophrenia and unipolar depression but not with bipolar disorder in Han Chinese. *Pharmacogenetics* 1999; **9**: 491–495.
- 92 Domotor E, Sarosi A, Balogh G, Szekeley A, Hejjas K, Sasvari-Szekely M *et al*. Association of neurocognitive endophenotype and STin2 polymorphism in major depressive disorder]. *Neuropsychopharmacol Hung* 2007; **9**: 53–62.
- 93 Bozina N, Mihaljevic-Peles A, Sagud M, Jakovljevic M, Sertic J. Serotonin transporter polymorphism in Croatian patients with major depressive disorder. *Psychiatr Danub* 2006; **18**: 83–89.
- 94 Battersby S, Ogilvie AD, Smith CA, Blackwood DH, Muir WJ, Quinn JP *et al*. Structure of a variable number tandem repeat of the serotonin transporter gene and association with affective disorder. *Psychiatr Genet* 1996; **6**: 177–181.
- 95 Gutierrez B, Pintor L, Gasto C, Rosa A, Bertranpetit J, Vieta E *et al*. Variability in the serotonin transporter gene and increased risk for major depression with melancholia. *Hum Genet* 1998; **103**: 319–322.
- 96 Schahab S, Heun R, Schmitz S, Maier W, Kolsch H. Association of polymorphism in the transcription factor LBP-1c/CP2/LSF gene with Alzheimer's disease and major depression. *Dement Geriatr Cogn Disord* 2006; **22**: 95–98.
- 97 Tan EC, Chan AO, Tan CH, Mahendran R, Wang A, Chua HC. Case-control and linkage disequilibrium studies of the tryptophan hydroxylase gene polymorphisms and major depressive disorder. *Psychiatr Genet* 2003; **13**: 151–154.
- 98 Gizatullin R, Zabolli G, Jonsson EG, Asberg M, Leopardi R. Haplotype analysis reveals tryptophan hydroxylase (TPH) 1 gene variants associated with major depression. *Biol Psychiatry* 2006; **59**: 295–300.
- 99 Zill P, Baghai TC, Zwanzger P, Schule C, Eser D, Rupprecht R *et al*. SNP and haplotype analysis of a novel tryptophan hydroxylase isoform (TPH2) gene provide evidence for association with major depression. *Mol Psychiatry* 2004; **9**: 1030–1036.
- 100 Zhou Z, Roy A, Lipsky R, Kuchipudi K, Zhu G, Taubman J *et al*. Haplotype-based linkage of tryptophan hydroxylase 2 to suicide attempt, major depression, and cerebrospinal fluid 5-hydroxyindoleacetic acid in 4 populations. *Arch Gen Psychiatry* 2005; **62**: 1109–1118.
- 101 Koido K, Koks S, Nikopensius T, Maron E, Altmae S, Heinaste E *et al*. Polymorphisms in wolframin (WFS1) gene are possibly related to increased risk for mood disorders. *Int J Neuropsychopharmacol* 2005; **8**: 235–244.